arXiv:math/0202091v4 [math.QA] 21 Dec 2007

DUALS OF POINTED HOPF ALGEBRAS

M. BEATTIE

Abstract. In this paper, we study the duals of some finite dimensional pointed Hopf algebras working over an algebraically closed field k of characteristic 0. In particular, we study pointed Hopf algebras with coradical $k[\Gamma]$ for Γ a finite abelian group, and with associated graded Hopf algebra of the form $\mathcal{B}(V)\#k[\Gamma]$ where $\mathcal{B}(V)$ is the Nichols algebra of $V = \oplus_i V_{g_i}^{\chi_i} \in {}^{k[\Gamma]}_{k[\Gamma]}\mathcal{YD}$. As a corollary to a general theorem on duals of coradically graded Hopf algebras, we have that the dual of $\mathcal{B}(V)\#k[\Gamma]$ is $\mathcal{B}(W)\#k[\hat{\Gamma}]$ where $W = \oplus_i W_{\chi_i}^{g_i} \in {}^{k[\hat{\Gamma}]}_{k[\hat{\Gamma}]}\mathcal{YD}$. This description of the dual is used to explicitly describe the Drinfel'd double of $\mathcal{B}(V)\#k[\Gamma]$. We also show that the dual of a nontrivial lifting A of $\mathcal{B}(V)\#k[\Gamma]$ which is not itself a Radford biproduct, is never pointed. For V a quantum linear space of dimension 1 or 2, we describe the duals of some liftings of $\mathcal{B}(V)\#k[\Gamma]$. We conclude with some examples where we determine all the irreducible finite-dimensional representations of a lifting of $\mathcal{B}(V)\#k[\Gamma]$ by computing the matrix coalgebras in the coradical of the dual.

1. Introduction and Preliminaries

Unless otherwise specified, we work over k, an algebraically closed field of characteristic 0. The object of this paper is to find the structure of dual Hopf algebras to some finite dimensional pointed Hopf algebras with coradical $k[\Gamma]$, with Γ a finite abelian group. A method for classifying pointed Hopf algebras has been proposed by N. Andruskiewitsch and H.-J. Schneider; an up-to-date survey of this project may be found in [4]. Roughly, this approach to studying pointed Hopf algebras is based on the following observations.

Let A be a Hopf algebra such that the coradical A_0 is a Hopf subalgebra of A so that the coradical filtration of A is a Hopf algebra filtration. Let $H = grA$ be the associated (coradically) graded Hopf algebra [34, Chapter XI], i.e., $H = \oplus_{n\geq 0} H(n)$ with $H(0) = A_0$ and $H(n) = A_n/A_{n-1}$ for $n \geq 1$. (Recall from [13] that a graded coalgebra $C = \oplus_{i=0}^{\infty} C(i)$ is coradically graded if $C(0) = C_0$ and $C_1 = C(0) \oplus C(1)$, where C_m denotes the coradical filtration and also if C_0 is one-dimensional then C is called strictly graded.) Then π, the Hopf algebra projection of H onto $H_0 = A_0$ [26, 5.4.2] splits the inclusion $i : H_0 \to H$ and thus $H \cong \mathcal{R}\#H_0$, the Radford biproduct of $\mathcal{R}$ and H_0, where $\mathcal{R} = H^{co\pi} = \{x \in H : (id \otimes \pi)\triangle(x) = x \otimes 1\}$ is the set of π-coinvariants. $\mathcal{R}$ is a Hopf algebra in ${}^{H_0}_{H_0}\mathcal{YD}$, the category of Yetter-Drinfel'd modules over H_0. In particular, if $H_0 = k[\Gamma]$, for Γ a finite abelian group, as will be the case in the examples of interest in this paper, then ${}^{k[\Gamma]}_{k[\Gamma]}\mathcal{YD}$ is the set of left $k[\Gamma]$-modules with a Γ-grading such that the Γ-action preserves the grading.

Research partially supported by NSERC.

Also $\mathcal{R}$ inherits a grading from H, i.e. $\mathcal{R}$ is graded by $\mathcal{R}(n) = \mathcal{R} \cap H(n)$ with $\mathcal{R}(0) = k \cdot 1$, and $\mathcal{R}(1) = P(\mathcal{R})$, the space of primitive elements of $\mathcal{R}$. If $V = \mathcal{R}(1)$ generates $\mathcal{R}$ as a Hopf algebra, then $\mathcal{R} = \mathcal{B}(V)$, the Nichols algebra of V. In any case $\mathcal{B}(V) \subseteq \mathcal{R}$. Much more detail about Nichols algebras can be found in [18] or in the original source material [29]. An excellent comprehensive survey of this approach with all the definitions and a compendium of results from various papers in the most generality may be found in [4].

Now, to construct a finite dimensional pointed Hopf algebra with coradical $k[\Gamma]$, the idea is to first find $V \in {}^{k[\Gamma]}_{k[\Gamma]}\mathcal{YD}$ with $\mathcal{B}(V)$ finite dimensional, and form the coradically graded Hopf algebra $H = \mathcal{B}(V)\#k[\Gamma]$. This construction is called bosonization or the Radford biproduct. A Hopf algebra A such that $grA \cong H$ is called a lifting of H and if $A \ncong grA$, then A is a nontrivial lifting of H. If V is a quantum linear space, or is of Cartan type A_n or of Cartan type B_2, the liftings of $\mathcal{B}(V)\#k[\Gamma]$ have been computed in [5], [9], [6], [4],[1], [10], respectively. We will study duals of liftings of some quantum linear spaces.

We abbreviate ${}^{k[\Gamma]}_{k[\Gamma]}\mathcal{YD}$ to ${}^{\Gamma}_{\Gamma}\mathcal{YD}$ for the category of Yetter-Drinfel'd modules over $k[\Gamma]$. For $V \in {}^{\Gamma}_{\Gamma}\mathcal{YD}$, we write V_g^{χ} with $g \in \Gamma, \chi \in \hat{\Gamma}$ for the set of $v \in V$ with the action of Γ on v given by $h \rightharpoonup v = \chi(h)v$ and the coaction by $\delta(v) = g \otimes v$. If Γ is an abelian group, and $V \in {}^{\Gamma}_{\Gamma}\mathcal{YD}$ then $V = \oplus_{\substack{g \in \Gamma \\ \chi \in \hat{\Gamma}}} V_g^{\chi}$ [4, Section 2]. If $V = \oplus_{i=1}^{t} kv_i$ with $v_i \in V_{g_i}^{\chi_i}$ then the braid matrix B of V is $(b_{ij}) = (\chi_j(g_i))$.

Under our assumptions on k and assuming Γ is finite abelian, we identify $k[\Gamma]^*$ with the Hopf algebra $k[\hat{\Gamma}]$ where $\hat{\Gamma}$ is the character group of Γ. Let $\varphi : k[\Gamma]^* \to k[\hat{\Gamma}]$ be the isomorphism between these Hopf algebras. For $g \in \Gamma$, the element $g^{**} \in Alg(k[\hat{\Gamma}], k)$ is defined by $g^{**}(\chi) = \chi(g)$. We usually simply write g for a character of $\hat{\Gamma}$ instead of g^{**}.

Unless otherwise noted, Γ will denote a finite abelian group.

For A a Hopf algebra with $g, h \in G(A)$, the group of grouplike elements of A, then $P(A)_{g,h} = \{x \in A | \triangle x = x \otimes g + h \otimes x\}$, the (g, h)-primitives of A. For each g, h, we write $P'(A)_{g,h}$ for a subspace of $P(A)_{g,h}$ such that $P(A)_{g,h} = k(g - h) \oplus P'(A)_{g,h}$ [26, §5.4].

We use the Sweedler summation notation throughout. $\overline{S}$ will denote the composition inverse to the antipode S.

Throughout, we write I_j for the $j \times j$ identity matrix. Also we use the notation $\mathcal{M}^c(r, k)$ for an $r \times r$ matrix coalgebra over k and we call a basis $e(i, j), 1 \leq i, j \leq r$, a matrix coalgebra basis if $\triangle(e(i, j)) = \sum_{k=1}^{r} e(i, k) \otimes e(k, j)$, and $\epsilon(e(i, j)) = \delta_{ij}$.

Our specific constructions and examples will focus on the case of V a quantum linear space.

Definition 1.1. *$V = \oplus_{i=1}^{t} kv_i \in {}^{\Gamma}_{\Gamma}\mathcal{YD}$ with $v_i \in V_{g_i}^{\chi_i}$ is called a quantum linear space if $\chi_i(g_j)\chi_j(g_i) = 1$ for $i \neq j$, i.e. in the braid matrix B, $b_{ij}b_{ji} = 1$ for $i \neq j$.*

Recall that for V a quantum linear space as above, $\mathcal{B}(V)\#k[\Gamma]$ is the Hopf algebra generated by the $(1, g_i)$-primitives $v_i\#1$ (usually written just v_i) and the grouplike elements $h \in \Gamma$. Multiplication is given by $hv_i = \chi_i(h)v_ih$ and $v_iv_j = \chi_j(g_i)v_jv_i$. Also if $\chi_i(g_i)$ is a primitive r_ith root of unity, $v_i^{r_i} = 0$, and then $\dim \mathcal{B}(V) = \Pi_{i=1}^t r_i$.

Proposition 1.2. *(see [5] or [9]) For V a quantum linear space with $\chi_i(g_i)$ a primitive r_ith root of unity, all liftings A of $\mathcal{B}(V)\#k[\Gamma]$ are Hopf algebras generated by the grouplikes and by $(1, g_i)$-primitives $x_i, 1 \le i \le t$ where*

$$hx_i = \chi_i(h)x_ih;$$

$$x_i^{r_i} = \alpha_{ii}(g_i^{r_i} - 1);$$

$$x_ix_j = \chi_j(g_i)x_jx_i + \alpha_{ij}(g_ig_j - 1).$$

We may assume $\alpha_{ii} \in \{0, 1\}$ and then we must have that

$$\alpha_{ii} = 0 \textit{ if } g_i^{r_i} = 1 \textit{ or } \chi_i^{r_i} \neq \epsilon \textit{ and } \alpha_{ij} = 0 \textit{ if } g_ig_j = 1 \textit{ or } \chi_i\chi_j \neq \epsilon. \tag{1.1}$$

Note that $\alpha_{ji} = -\chi_j(g_i)^{-1}\alpha_{ij} = -\chi_i(g_j)\alpha_{ij}$. Thus the lifting A is described by a matrix $\mathcal{A} = (\alpha_{ij})$ with 0's or 1's down the diagonal and with $\alpha_{ji} = -\chi_i(g_j)\alpha_{ij}$ for $i \neq j$. We associate $\mathcal{A} = (0)$ with $\mathcal{B}(V)\#k[\Gamma]$. □

2. The dual of a Radford biproduct.

The first theorem in this section is in the form suggested by N. Andruskiewitsch who points out that the ideas for this proof are already in the literature although the statement does not appear explicitly. (See [7], [3], [29].) The theorem in the original version of this paper corresponded to Corollary 2.3 and follows directly from this more general formulation. In fact it is shown in [3, 3.2.30] that for $V \in {}_H^H\mathcal{YD}$ with $\mathcal{B}(V)$ finite dimensional, then $\mathcal{B}({}^*V) \cong ({}^*\mathcal{B}(V))^{opcop}$ in the category ${}_H^H\mathcal{YD}$. Note that the categories ${}_H^H\mathcal{YD}$ and ${}_{H^*}^{H^*}\mathcal{YD}$ are equivalent for H finite dimensional with bijective antipode by [3, 2.2.1]. The results of Corollary 2.3 give a special case.

For k any field, let H be a graded Hopf algebra, $H = \oplus_{n\ge 0}H^n$ with finite dimensional components H^n. Let $L = \oplus_{n\ge 0}(H^n)^*$, with the "dual" Hopf algebra structure. Assume that $H_0 = H^0$, i.e., H^0 is cosemisimple. (Note that without restriction on k this does not imply that L^0 is cosemisimple.) Let $\pi : H \to H^0$, $p : H \to H^1$, $\sigma : L \to L^0$ be the homogeneous projections. Then, as discussed in Section 1, $H \simeq \mathcal{R}\#H^0$, and $L \simeq \mathcal{S}\#L^0$ where $\mathcal{R} = H^{co\pi}$ and $\mathcal{S} = L^{co\sigma}$.

Lemma 2.1. *For k any field and H, L as above, the following are equivalent:*

(i) H is coradically graded.

(ii) L is generated as an algebra by L^0 and L^1.

Proof. Suppose (i) holds. By [13, 2.2], H is coradically graded if and only if $H_1 = H^0 + H^1$. Now consider the map ϕ from H^2 to $H^1 \otimes H^1$ defined by $\phi := p \otimes p \circ \triangle$. If $\phi(a) = 0$, then $\triangle(a) \in H^0 \otimes H^2 + H^2 \otimes H^0$, and then since $H^0 = H_0$, $a \in H_1$, which is impossible unless $a = 0$ since $H_1 \cap H^2 = 0$. Thus ϕ is injective. By induction (or see the proof of [7, Lemma 5.5]), the map $(p \otimes p \otimes \ldots \otimes p) \circ \triangle^{n-1}$ from H^n, the nth homogeneous component of H, to $H^1 \otimes \ldots \otimes H^1$ is injective for $n \ge 2$. Clearly this is equivalent to the surjectivity of the multiplication map from $L^1 \otimes L^1 \otimes \ldots \otimes L^1$ to L^n.

Conversely, if the multiplication map from $L^1 \otimes L^1 \otimes \ldots \otimes L^1$ to L^n is surjective, so that the map $(p \otimes p \otimes \ldots \otimes p) \circ \triangle^{n-1}$ from H^n to $H^1 \otimes \ldots \otimes H^1$ is injective, then since $(p \otimes p \otimes \ldots \otimes p) \circ \triangle^{n-1}$ acting on H_1 is zero for any $n > 1$, then $H_1 \cap H^n = 0$ for all $n \geq 2$. Thus, the containment $H^1 + H^0 \subseteq H_1$ is an equality, and H is coradically graded. □

From the lemma above, we immediately obtain:

Theorem 2.2. *Let k be any field and let H and L be as above, with the additional assumption that $H^0 = H_0$ is cosemisimple and also semisimple, so that L_0 is also. The following are equivalent:*

(i) H is coradically graded and is generated as an algebra by H^0 and H^1.

(ii) L is coradically graded and is generated as an algebra by L^0 and L^1.

A Hopf algebra H satisfying the conditions of Theorem 2.2 is called a "bialgebra of type one" in the original terminology of Nichols' [29]. Then, as briefly outlined in Section 1 and explained in detail in [5, Lemmas 2.4, 2.5] or in the survey [4, Section 2], $H \simeq \mathcal{B}(V)\#H^0$, where $V = \mathcal{R}(1) = P(\mathcal{R})$, the space of primitives of $\mathcal{R} = \mathcal{B}(V)$, and $L \simeq \mathcal{B}(W)\#L^0$, where $W = \mathcal{S}(1) = P(\mathcal{S})$. Here $\mathcal{R} = \mathcal{B}(V)$ and $\mathcal{S} = \mathcal{B}(W)$ are Nichols algebras; in particular, they are strictly graded coradically graded Hopf algebras in the categories ${}^{H_0}_{H_0}\mathcal{YD}$ and ${}^{L_0}_{L_0}\mathcal{YD}$ respectively and are generated as algebras by their 1-graded component.

There is a relation between the Yetter-Drinfeld module structures on V and W above. Observe that the pairing $H^1 \times L^1 \to k$ gives rise to a pairing $(\ ,\ ) : V\#H^0 \times W\#L^0 \to k$, which is explicitly $(v\#x, w\#y) = (v, w)(x, y)$ (by the definition of $\mathcal{R}$ and $\mathcal{S}$). Next, if $\delta : W \to L^0 \otimes W$ is the coaction, consider the right action $\leftharpoonup : W \otimes H^0 \to W$, $w \leftharpoonup a := (a \otimes Id)\delta(w)$. Then since, by the definition of $W \subset \mathcal{S} = L^{co\sigma}$, for $w \in W$, we have $\sum w_{(1)} \otimes w_{(2)} \otimes \sigma(w_{(3)}) = \sum w_{(1)} \otimes w_{(2)} \otimes 1$, and then

$$(ad(a)v, w) = (v, w \leftharpoonup a), \qquad a \in H^0, \tag{2.2}$$

where $ad(a)v$ denotes the adjoint action, i.e. $ad(a)v = \sum a_1 v S(a_2)$. Interchanging rôles, we obtain the relation between the coaction of V and the action of W.

Now we resume our assumption that k is algebraically closed and of characteristic 0, and Γ is a finite abelian group. Let $V \in {}^{\Gamma}_{\Gamma}\mathcal{YD}$ with basis $\{v_1, \ldots, v_t\}$ where $v_i \in V^{\chi_i}_{g_i}, g_i \in \Gamma, \chi_i \in \hat{\Gamma}$. Suppose $\mathcal{B}(V)$, the Nichols algebra, is a finite dimensional Hopf algebra in ${}^{\Gamma}_{\Gamma}\mathcal{YD}$. Assume $\chi_i(g_i)$ is a primitive r_ith root of unity for some integer $r_i > 1$, and then $v_i^{r_i} = 0$ in $\mathcal{B}(V)$.

Corollary 2.3. *Let $H = \mathcal{B}(V)\#k[\Gamma]$ for k, V, Γ as above. Then $H^* \cong \mathcal{B}(W)\#k[\hat{\Gamma}]$ where $W = \oplus_{i=1}^{t} k w_i \in {}^{\hat{\Gamma}}_{\hat{\Gamma}}\mathcal{YD}$ and $w_i \in W^{g_i}_{\chi_i}$.*

Proof. Let $w_i \in W$ be defined by $(v_j, w_i) = \delta_{i,j}$. Suppose $\delta(w_j) = \sum_{k=1}^{t} \varphi_{jk} \otimes w_k \in k[\hat{\Gamma}] \otimes W$. Then for $\leftharpoonup$ the right action of Γ on W described above, $w_j \leftharpoonup h = \sum_{k=1}^{t} \varphi_{jk}(h) w_k$ and so by (2.2), $\varphi_{jk}(h) = (v_k, w_j \leftharpoonup h) = (ad(h)v_k, w_j) =$

$\delta_{kj}\chi_k(h)$. Thus $\delta(w_j) = \chi_j \otimes w_j$. A similar argument shows that for $\chi \in \hat{\Gamma}, \chi \cdot w_i = \chi(g_i)w_i$. □

Recall from Section 1, that a pointed Hopf algebra A is called a lifting of $\mathcal{B}(V)\#k[\Gamma]$ if grA, the associated graded Hopf algebra, is isomorphic to $\mathcal{B}(V)\#k[\Gamma]$. The lifting is nontrivial if $A \ncong grA$. Now we show that, although the dual of $H = \mathcal{B}(V)\#k[\Gamma]$ is pointed, the dual of a nontrivial lifting A of H is not pointed if the coalgebra projection π from A to A_0 is not an algebra map. This result has been shown for various particular cases in the literature, for example, [32, p.315], [17, Theorem 2.5], [15, Cor.5.3].

Proposition 2.4. *Let A be a nontrivial lifting of $H = \mathcal{B}(V)\#k[\Gamma]$ such that $\pi : A \longrightarrow A_0$ is not an algebra map. Then $G(A^*)$, the group of grouplikes of A^*, is isomorphic to a proper subgroup of $\hat{\Gamma}$ and A^* is not pointed.*

Proof. As above, since A is a lifting of H, A is generated as an algebra by $(1, g_i)$-primitives x_i with $x_i \in P'_{(1,g_i)}$ and the elements of Γ. The x_i are the liftings of the $v_i\#1 \in \mathcal{B}(V)\#k[\Gamma]$. Also $g_i x_i = \chi_i(g_i)x_i g_i$ and since $\chi_i(g_i) \neq 1$, all grouplike elements of A^* map the x_i to 0. Thus grouplike elements of A^* are nonzero only on $A_0 = k[\Gamma]$ and so all grouplikes are of the form $\gamma \cdot \pi$ where $\gamma \in \hat{\Gamma}$.

However not every $\gamma \in \hat{\Gamma}$ is grouplike in A^*. Since π is not an algebra map, there exist $x, y \in A$ such that $\pi(x)\pi(y) \neq \pi(xy)$ in $k[\Gamma]$. But then there must exist $\gamma \in \hat{\Gamma}$ such that $\gamma(\pi(x)\pi(y)) = \gamma\pi(x)\gamma\pi(y) \neq \gamma\pi(xy)$, for otherwise $\varphi(\pi(x)\pi(y)) = \varphi(\pi(xy))$ for all $\varphi \in k[\Gamma]^*$. Then $\gamma\pi$ is not an algebra map from A to k and $\gamma \notin G(A^*)$. Thus $G(A^*)$ is a proper subgroup of $\hat{\Gamma}$.

Let $Jac(\cdot)$ denote the Jacobson radical. Since $Jac(k[\Gamma]) = 0$ and $k[\Gamma]$ is commutative, we have that $k[\Gamma] \cap Jac(A) = 0$. For if $z \in k[\Gamma] \cap Jac(A)$, then z is nilpotent by a theorem of Amitsur [22, 4.20] and so lies in $Jac(k[\Gamma])$. Thus dim Jac$(A) \leq (\dim A) - |\Gamma|$ as k-spaces, and by [26, 5.1.7], $\dim corad(A^*) = \dim(A) - \dim (\text{Jac}(A)) \geq |\Gamma|$. Thus A^* cannot be pointed. □

Thus the coradical of the dual of a nontrivial lifting of a quantum linear space contains nontrivial matrix coalgebras. In the next sections, we will discuss the number and dimensions of these matrix coalgebras for some particular examples.

We now use Corollary 2.3 to compute the Drinfel'd double $D(H)$ for a finite dimensional Hopf algebra $H = \mathcal{B}(V)\#k[\Gamma]$. Suppose, as above, that $V = \oplus_{i=1}^t kv_i$ where $v_i \in V_{g_i}^{\chi_i}$. Also, as above, define $w_i \in W_{\chi_i}^{g_i} \subset H^*$ by $(v_j, w_i) = \delta_{i,j}$. Then $w_i(hv_i) = \chi_i(h)$ and $w_i \in P(H^*)_{\epsilon,\chi_i}$. Define $w_i' = \chi_i^{-1} * w_i = (1\#\chi_i^{-1})(w_i\#\epsilon)$ in H^* . Then $w_i'(z) = \chi_i^{-1}(g_i)$ for z in $H(1)$ of the form hv_i and 0 otherwise. In H^*, w_i' is $(\chi_i^{-1}, \epsilon_H)$ -primitive. Let $W' = \oplus_{i=1}^t kw_i'$.

As usual, the left (right) action of H on H^* is denoted $h \rightharpoonup h^*$ $(h^* \leftharpoonup h)$ where $h \rightharpoonup h^*(z) = h^*(zh)$ $(h^* \leftharpoonup h(z) = h^*(hz))$.

Theorem 2.5. *Let $H = \mathcal{B}(V)\#k[\Gamma]$, with $V = \oplus_{i=1}^t kv_i$ and $v_i \in V_{g_i}^{\chi_i}$. Let $D(H)$ be the Drinfel'd double of H. Then $gr(D(H)) \simeq \mathcal{B}(U)\#k[\hat{\Gamma} \times \Gamma]$ where $U \in {}_{\hat{\Gamma}\times\Gamma}^{\hat{\Gamma}\times\Gamma}\mathcal{YD}$ is the direct sum of W' and V with appropriate actions and coactions, i.e. $U =$*

$\oplus_{i=1}^{2t} k u_i$ where $u_i \in U^{g_i \times \chi_i^{-1}}_{\chi_i^{-1} \times 1}$ for $i = 1, \dots, t$ and $u_i \in U^{g_i^{-1} \times \chi_i}_{\epsilon \times g_i}$ for $i = t+1, \dots, 2t$. Also $D(H) \ncong gr(D(H))$.

Proof. In H^{*cop}, the element w_i' is $(\epsilon_H, \chi_i^{-1})$-primitive and $\gamma w_i' \gamma^{-1} = g_i^{**}(\gamma) w_i'$ for $\gamma \in G(H^{*cop})$ so that $w_i' \in P(H^{*cop})^{g_i}_{\chi_i^{-1}}$. Thus by an argument similar to that in Corollary 2.3, $H^{*cop} \cong \mathcal{B}(W')\#k[\hat{\Gamma}]$. Since, as coalgebras, $D(H) = H^{*cop} \otimes H$, by [26, 5.1.10] or [32], $D(H)$ is pointed with group of grouplikes $\hat{\Gamma} \times \Gamma$.

In $D(H) = H^{*cop} \bowtie H$, the elements $\epsilon \bowtie v_i$ with $v_i \in V$ are $(\epsilon \bowtie 1, \epsilon \bowtie g_i)$-primitive and the $w_i' \bowtie 1$ with $w_i' \in W'$ are $(\epsilon \bowtie 1, \chi_i^{-1} \bowtie 1)$-primitive. For $\gamma \bowtie h \in G(D(H))$, using the formula for multiplication in $D(H)$ given in [26, 10.3.11 (1)] from [32] , we compute

$$\begin{aligned}(\gamma \bowtie h)(\epsilon \bowtie v_i)(\gamma^{-1} \bowtie h^{-1}) &= (\gamma \bowtie h v_i)(\gamma^{-1} \bowtie h^{-1}) \\ &= \sum \gamma[(hv_i)_1 \rightharpoonup \gamma^{-1} \leftharpoonup \overline{S}((hv_i)_3)] \bowtie (hv_i)_2 h^{-1}\end{aligned}$$

where the left and right actions of H on H^* are the usual ones. Now, $\triangle^2(hv_i) = hv_i \otimes h \otimes h + hg_i \otimes hv_i \otimes h + hg_i \otimes hg_i \otimes hv_i$ and $\overline{S}(hv_i) = S^{2r_i - 1}(hv_i) = -\chi_i(g_i h)(g_i h)^{-1} v_i$. Since $hv_i \rightharpoonup \gamma^{-1}$ and $\gamma^{-1} \leftharpoonup l v_i$ are 0 for any $h, l \in \Gamma$, then

$$\begin{aligned}\sum \gamma[(hv_i)_1 \rightharpoonup \gamma^{-1} \leftharpoonup \overline{S}((hv_i)_3)] \bowtie (hv_i)_2 h^{-1} &= \gamma[hg_i \rightharpoonup \gamma^{-1} \leftharpoonup h^{-1}] \bowtie hv_i h^{-1} \\ = \gamma^{-1}(g_i)\epsilon \bowtie \chi_i(h) v_i &= (g_i^{**-1} \otimes \chi_i)(\gamma \otimes h)\epsilon \bowtie v_i.\end{aligned}$$

Then $\epsilon \bowtie v_i \in P(D(H))^{g_i^{-1} \bowtie \chi_i}_{\epsilon \bowtie g_i}$ where g_i^{-1} means the usual identification of $Alg(\hat{\Gamma}, k)$ with Γ. Similarly, we see that $w_i' \bowtie 1_H \in P(D(H))^{g_i \bowtie \chi_i^{-1}}_{\chi_i^{-1} \bowtie 1}$. Thus $U = W' \oplus V \in^{\hat{\Gamma} \times \Gamma}_{\hat{\Gamma} \times \Gamma} \mathcal{YD}$ with actions and coactions as in the statement of the theorem.

Now, $grD(H) \cong \mathcal{R}\#k[\hat{\Gamma} \times \Gamma]$ as Hopf algebras. Then $\mathcal{B}(W' \oplus V) \subseteq \mathcal{R}$. If $1 \leq i, j \leq t$, in the braid matrix for $W' \oplus V$, $b_{i,t+j} =< g_j^{-1} \bowtie \chi_j, \chi_i^{-1} \bowtie 1 >= \chi_i(g_j)$ and $b_{t+j,i} =< g_i \bowtie \chi_i^{-1}, \epsilon_H \bowtie g_j >= \chi_i(g_j)^{-1}$. Thus by [18, Theorem 2.2], $\dim \mathcal{B}(W' \oplus V) = \dim \mathcal{B}(W') \dim \mathcal{B}(V) = \dim \mathcal{R}$. Thus $grD(H) \cong \mathcal{B}(W' \oplus V)\#k[\hat{\Gamma} \times \Gamma]$.

To show that $D(H) \ncong grD(H)$, we compute the products of the v_i and w_j'. By [26, 5.1.10], $(\epsilon_H \bowtie v_i)(w_j' \bowtie 1_H) = \sum v_{i_1} \rightharpoonup w_j' \leftharpoonup \overline{S}(v_{i_3}) \bowtie v_{i_2} = v_i \rightharpoonup w_j' \bowtie 1 + g_i \rightharpoonup w_j' \bowtie v_i + g_i \rightharpoonup w_j' \leftharpoonup \overline{S}(v_i) \bowtie g_i$. However $v_i \rightharpoonup w_j'$ and $w_j' \leftharpoonup \overline{S}(v_i)$ are 0 if $i \neq j$ and then $v_i w_j' = \chi_j^{-1}(g_i) w_j' v_i$. If $i = j$, then $v_i \rightharpoonup w_i' = \chi_i(g_i)^{-1}\epsilon$ and $g_i \rightharpoonup w_i' \leftharpoonup \overline{S}(v_i) = -\chi_i(g_i)^{-1}\chi_i$, so that

$$v_i w_i' = \chi_i(g_i)^{-1} w_i' v_i + \chi_i(g_i)^{-1}(\epsilon \bowtie 1 - \chi_i \bowtie g_i).$$

Thus π is not an algebra map and $D(H) \ncong \mathcal{B}(U)\#k[\hat{\Gamma} \times \Gamma]$ as algebras. □

Note that by adjusting the skew-primitives by a scalar in the above theorem, we may assume that $u_{t+j} u_j = \chi_j^{-1}(g_j) u_j u_{t+j} + (\chi_j^{-1} \bowtie g_j - \epsilon \bowtie 1)$. Recall from [32] that $D(H)$ is always minimal quasitriangular and that the dual $D(H)^*$ is also quasitriangular if and only if H and H^* are quasi-triangular.

For V a quantum linear space, the question of when Hopf algebras $\mathcal{B}(V)\#k[\Gamma]$, their liftings and their doubles are quasi-triangular or ribbon has been considered in [21], [33], [17], [30], [27], [28].

Example 2.6. If V is a quantum linear space, then note that for $V, W' \in^{\hat{\Gamma}\times\Gamma}_{\hat{\Gamma}\times\Gamma} \mathcal{YD}$ as in Theorem 2.5, $U = W' \oplus V \in^{\hat{\Gamma}\times\Gamma}_{\hat{\Gamma}\times\Gamma} \mathcal{YD}$ is also a quantum linear space. Thus if $H = \mathcal{B}(V)\#k[\Gamma]$, $A = D(H)$ is the nontrivial lifting of $\mathcal{B}(W' \oplus V)\#k[\hat{\Gamma} \times \Gamma]$ with lifting matrix $\mathcal{A} = \begin{bmatrix} 0 & D \\ I_t & 0 \end{bmatrix}$ where D is the $t \times t$ diagonal matrix with $d_{ii} = -\chi_i(g_i)$. □

The quasi-triangularity of $D(H)$ can be used to prove the quasi-triangularity of some Radford biproducts.

Example 2.7. Let $V, W', U, H, A = D(H)$ be as in Example 2.6. Furthermore, let $\chi_i(g_i) = -1$ for all $1 \leq i \leq t$. Let J be the Hopf ideal of A generated by the skew-primitives $\chi_i^{-1} \bowtie g_i - \epsilon \bowtie 1$ where $1 \leq i \leq t$. Let $\Gamma' = (\hat{\Gamma} \times \Gamma)/\Lambda$ where Λ is the subgroup generated by the $\chi_i^{-1} \bowtie g_i$. Then U has a natural Γ' grading and also a Γ'-action since because $\chi_i(g_i)^2 = 1 = \chi_i(g_j)\chi_j(g_i)$ then $< g_j^{-1} \bowtie \chi_j, \chi_i^{-1} \bowtie g_i > = < g_j \bowtie \chi_j^{-1}, \chi_i^{-1} \bowtie g_i > = 1$ for all $1 \leq i, j \leq t$. In fact, it is easy to see that $A/J \cong \mathcal{B}(U)\#k[\Gamma']$ where U has this Γ'-action and grading. Thus this biproduct is minimal quasi triangular since A is. The R-matrices on $L = \mathcal{B}(U)\#k[\Gamma']$ are discussed in [28], as well as the existence of ribbon elements for L and $D(L)$. □

If $\Gamma \cong C_2$, then $\mathcal{B}(U)\#k[\Gamma'] = E(2t)$ and this Hopf algebra has been studied by many authors.

Example 2.8. Let U be the quantum linear space $U = \oplus_{i=1}^{2t} kv_i \in^{\hat{\Gamma}\times\Gamma}_{\hat{\Gamma}\times\Gamma} \mathcal{YD}$ with $\Gamma =< g >\cong C_2$ and $\hat{\Gamma} =< \chi >\cong C_2$ with $v_i \in U^{(g,\chi)}_{(1,g)}$ for $1 \leq i \leq t$, $v_j \in U^{(g,\chi)}_{(\chi,1)}$ for $t+1 \leq j \leq 2t$. Let A be the nontrivial lifting of $\mathcal{B}(U)\#k[\hat{\Gamma} \times \Gamma]$ with commutation matrix $\begin{bmatrix} 0 & I_t \\ I_t & 0 \end{bmatrix}$. Then A is minimal quasitriangular since for $V = \oplus_{i=1}^{t} kv_i$, we have $A = D(\mathcal{B}(V)\#k[\Gamma]) = D(E(t))$ in the notation of [30]. Also it is shown in [30] that $A = D(E(t))$ has quasi-ribbon elements if and only if t is even, in which case it has 4 quasi-ribbon elements, two of which are ribbon. And, since it is well-known that $E(t)$ is quasitriangular and by [9] or by Corollary 2.3, $E(t)$ is self-dual, then $A^* = D(E(t))^*$ is also quasitriangular, but is not pointed by Proposition 2.4. If $t = 1$, then A is the Drinfel'd double of the Sweedler Hopf algebra. This 16 dimensional pointed Hopf algebra is discussed in [8, Section 2, Case 2] and in [11, p.562] as $H_{(5)}$, and in [32, p.315], where the fact that A^* is not pointed is proved directly. The dual of $D(E(1))$ will be computed in Section 4. □

Remark 2.9. N. Andruskiewitsch has also pointed out that Theorem 2.5 has an antecedent in Drinfel'd's paper [14] where $U_q(g)$ was introduced as a quotient of the quantum double of $U_q(b)$, and that this material is strongly tied to the construction in [23] where Lusztig defines $U_q(g)$ via what are now known as Nichols algebras. Theorem 2.5 can also be obtained as a corollary to a construction of a (possibly infinite) Hopf algebra $\mathcal{D}(V)$ using the notation and arguments of [23].

Here is an outline of the proof. For $V, W, \Gamma, \hat{\Gamma}$ as above, define a bilinear form from $(\mathcal{B}(W)\#k[\hat{\Gamma}]) \times (\mathcal{B}(V)\#k[\Gamma])$ to k by $(\xi\#z|x\#g) = (\xi, x)(z, g)$. Following [24, 7.2.6], form the graded double $\mathcal{D}(V)$. Identify V (respectively W) with subspaces of $\mathcal{D}(V)$ and determine the commutation relations for $v_i w_j$, $g w_j$, γv_i, and $g\gamma$ where $g \in \Gamma, \gamma \in \hat{\Gamma}$ as was done in the proof of Theorem 2.5. Then $\mathcal{D}(V)$ is isomorphic to $T(W \otimes k[\hat{\Gamma}] \otimes V \otimes k[\Gamma])/\mathcal{J}$ for $\mathcal{J}$ the two-sided ideal generated by the commutation relations.

As well, the coradical filtration of $\mathcal{D}(V)$ is given by

$$\begin{aligned}
&\mathcal{D}(V)_0 = k[\hat{\Gamma} \times \hat{\Gamma}];\\
&\mathcal{D}(V)_1 = \mathcal{D}(V)_0 + (W\#k[\hat{\Gamma}]) \bowtie k[\Gamma] + k[\hat{\Gamma}] \bowtie (V\#k[\Gamma]);\\
&\mathcal{D}(V)_n = (\mathcal{D}(V)_1)^n \text{ for } n > 1.
\end{aligned}$$

□

3. Duals of liftings of quantum linear spaces

Throughout this section let $V = \oplus_{i=1}^{t} k v_i \in {}^{\Gamma}_{\Gamma}\mathcal{YD}$ be a quantum linear space. We study duals of Hopf algebras A where A is a nontrivial lifting of $\mathcal{B}(V)\#k[\Gamma]$ with matrix $\mathcal{A}$ as in Proposition 1.2. For $v_i \in V_{g_i}^{\chi_i}$, suppose $\chi_i(g_i)$ is a primitive r_i-th root of unity. Let $x_i \in A$ be the lifting of $v_i\#1$ to A, i.e. in A, x_i is $(1, g_i)$-primitive and $hx_i = \chi_i(h)x_i h$ for all $h \in \Gamma$. Then A has a vector space basis hz with $h \in \Gamma$ and $z \in \mathcal{Z} = \{x_1^{m_1} \dots x_t^{m_t} | 0 \leq m_i \leq r_i - 1\}$.

For $\gamma \in \hat{\Gamma}$, define $w(\gamma, x_1^{m_1} \dots x_t^{m_t}) \in A^*$ to be the map which takes $hx_1^{m_1} \dots x_t^{m_t}$ to $\gamma(h)$ and all other basis elements to 0, so that $\mathcal{W} = \{w(\gamma, z) | \gamma \in \hat{\Gamma}, z \in \mathcal{Z}\}$ is a vector space basis for A^*. As in Section 2, we denote $w(\chi_i, x_i)$ by w_i and also $w(\gamma, 1)$ is usually just written γ.

Lemma 3.1. *Let A be a lifting of $\mathcal{B}(V)\#k[\Gamma]$ with V a quantum linear space. Let $\gamma, \lambda \in \hat{\Gamma}$, $0 \leq m_i \leq r_i - 1$ and $1 \leq i < j \leq t$.*

(i) Then

$$w(\gamma, x_i^{m_i}) * w(\lambda, x_j^{m_j}) = (\gamma\chi_i^{-m_i})(g_j^{m_j}) w(\gamma\lambda, x_i^{m_i} x_j^{m_j});$$
$$w(\lambda, x_j^{m_j}) * w(\gamma, x_i^{m_i}) = \lambda(g_i^{m_i}) w(\gamma\lambda, x_i^{m_i} x_j^{m_j}).$$

In particular, $w_i w_j = w(\chi_i\chi_j, x_i x_j) = \chi_i(g_j) w_j w_i$ for $i < j$.

ii)

$$\gamma * w(\lambda, x_1^{m_1} \dots x_t^{m_t}) = \gamma(g_1^{m_1} \dots g_t^{m_t}) w(\lambda\gamma, x_1^{m_1} \dots x_t^{m_t});$$
$$w(\lambda, x_1^{m_1} \dots x_t^{m_t}) * \gamma = w(\gamma\lambda, x_1^{m_1} \dots x_t^{m_t}).$$

In particular $\gamma * w_i = \gamma(g_i) w_i * \gamma = \gamma(g_i) w(\chi_i\gamma, x_i)$.

iii) For $0 \leq m_i \leq r_i - 1$ we have that

$$w(\lambda, x_i) * w(\gamma, x_i^{m_i}) = \lambda(g_i^{m_i})(q^{m_i} + q^{m_i - 1} + \dots 1) w(\lambda\gamma, x_i^{m_i + 1})$$

where $q = \chi_i(g_i)^{-1}$, a primitive r_ith root of 1.

The multiplication formulas above show that $\gamma \in \hat{\Gamma}$ and the elements w_i generate A^* as an algebra with $\gamma * w_i = \gamma(g_i) w_i * \gamma$, $w_i * w_j = \chi_i(g_j) w_j * w_i$ for $i \neq j$, and $w_i^{r_i} = 0$.

Proof. We prove i). The proofs of ii) and iii) are similar and straightforward. The map $w(\gamma, x_i^{m_i}) \otimes w(\lambda, x_j^{m_j})$ is nonzero only on elements of $A \otimes A$ of the form $hx_i^{m_i} \otimes lx_j^{m_j}$. Thus the only basis element on which $w(\gamma, x_i^{m_i}) * w(\lambda, x_j^{m_j})$ or $w(\lambda, x_j^{m_j}) * w(\gamma, x_i^{m_i})$ is nonzero is $hx_i^{m_i}x_j^{m_j}$. Then we have that

$$\begin{aligned} w(\gamma, x_i^{m_i}) * w(\lambda, x_j^{m_j})(hx_i^{m_i}x_j^{m_j}) &= w(\gamma, x_i^{m_i})(hx_i^{m_i}g_j^{m_j})w(\lambda, x_j^{m_j})(hx_j^{m_j}) \\ &= (\gamma\chi_i^{-m_i})(g_j^{m_j})\gamma\lambda(h). \end{aligned}$$

Also

$$\begin{aligned} w(\lambda, x_j^{m_j}) * w(\gamma, x_i^{m_i})(hx_i^{m_i}x_j^{m_j}) &= w(\lambda, x_j^{m_j})(hg_i^{m_i}x_j^{m_j})w(\gamma, x_i^{m_i})(hx_i^{m_i}) \\ &= (\lambda\gamma)(h)\lambda(g_i^{m_i}), \end{aligned}$$

which proves i). □

Proposition 3.2. *Let $V = \oplus_{i=1}^t kv_i$ be a quantum linear space, and let A be a lifting of $\mathcal{B}(V)\#k[\Gamma]$ with matrix $\mathcal{A}$ such that for $i \neq j$, if $\alpha_{i,j} \neq 0$, then $\chi_i = \chi_j$ so that $\chi_i^2 = \chi_j^2 = \epsilon$ and $r_i = r_j = 2$. We make the following definitions:*

Θ**:** *is the subgroup of $\hat{\Gamma}$ generated by the $\chi_i, 1 \leq i \leq t$;*
Ω**:** *is the subgroup of $\hat{\Gamma}$ defined by $\Omega = \{\gamma \in \hat{\Gamma} : \gamma(\alpha_{i,i}(g_i^{r_i} - 1)) = 0 = \gamma(\alpha_{i,j}(g_ig_j - 1))$ for $1 \leq i, j \leq t, i \neq j\}$;*
Γ'**:** *is the subgroup of Γ generated by $\{g_i^{r_i} | \alpha_{i,i} \neq 0\} \cup \{g_ig_j | \alpha_{i,j} \neq 0\}$.*

Let $\overline{\Gamma} = \Gamma/\Gamma'$ and note that $\hat{\overline{\Gamma}} \cong \Omega$. For $g \in \Gamma$, $\overline{g}$ will denote its image in $\overline{\Gamma}$. For $\gamma \in \hat{\Gamma}$, $\overline{\gamma}$ will denote its image in $\hat{\Gamma}/\Theta$. Then

1. *Θ is a subgroup of Ω and Ω is the group of grouplikes in A^*.*
2. *There exist subHopf algebras of A^* isomorphic to $\mathcal{B}(W)\#k[\Theta]$ and $\mathcal{B}(W)\#k[\Omega]$ where $W = \oplus_{i=1}^t kw_i$ with $w_i \in W_{\chi_i}^{\overline{g_i}}$. Denote by $C(\overline{1})$ the pointed subHopf algebra isomorphic to $\mathcal{B}(W)\#k[\Theta]$ and by B the subHopf algebra isomorphic to $\mathcal{B}(W)\#k[\Omega]$.*
3. *For $\gamma \in \hat{\Gamma}$, let $C(\overline{\gamma}) = \gamma C(\overline{1}) = C(\overline{1})\gamma$ for any $\gamma \in \overline{\gamma}$. $C(\overline{\gamma})$ is a subcoalgebra of A^* and as a coalgebra, $A^* \cong \oplus_{\overline{\gamma} \in \hat{\Gamma}/\Theta} C(\overline{\gamma})$. If $\gamma \in \Omega$, then $C(\overline{\gamma}) \cong C(\overline{1})$ as a coalgebra and $B \cong \oplus_{\overline{\gamma} \in \Omega/\Theta} C(\overline{\gamma})$ as a coalgebra. Thus $coradA^* \cong k[\Omega] \oplus \oplus_{\substack{\overline{\gamma} \in \hat{\Gamma}/\Theta \\ \gamma \notin \Omega}} (coradC(\overline{\gamma}))$.*
4. *For all $\overline{\gamma}, \overline{\lambda} \in \hat{\Gamma}/\Theta$, $C(\overline{\gamma})C(\overline{\lambda}) = C(\overline{\gamma\lambda})$ and also $S(C(\overline{\gamma})) = C(\overline{\gamma^{-1}})$.*

Proof. (1) Clearly $\gamma \in \hat{\Gamma}$ is grouplike in A^* if and only if $\gamma \in \Omega$. Now, if $\alpha_{i,i} \neq 0$ then $\chi_i^{r_i} = \epsilon$ and thus for $j \neq i$, $\chi_j(g_i^{r_i}) = \chi_i^{r_i}(g_j^{-1}) = 1$ so $\chi_j(\alpha_{i,i}(g_i^{r_i} - 1)) = 0$ for all i. Also, if $\alpha_{i,j} \neq 0$, then for $k \neq i, j$, $\chi_k(g_ig_j) = (\chi_i\chi_j)(g_k) = 1$. As well, $\chi_i(g_ig_j) = \chi_i(g_i)\chi_i(g_j) = \chi_i(g_i)\chi_j^{-1}(g_i) = (\chi_i\chi_j)(g_i) = 1$ since we are assuming here that $\chi_j^2 = \epsilon$.

(2) Let $\overline{V} = \oplus_{i=1}^t k\overline{v_i} \in {}^{\overline{\Gamma}}_{\overline{\Gamma}}\mathcal{Y}D$ with $\delta(\overline{v_i}) = \overline{g_i} \otimes \overline{v_i}, \overline{h} \rightarrow \overline{v_i} = \overline{h \rightarrow v_i} = \chi_i(h)\overline{v_i}$ where $\overline{g_i}, \overline{h}$, denote the images of g, h in $\overline{\Gamma}$. The action of $\overline{\Gamma}$ on $\overline{V}$ is well-defined by the arguments in (1). There is an exact sequence of Hopf algebras,

$$\begin{tikzcd} 1 \arrow[r] & k[\Gamma'] \arrow[r] & A \arrow[r, "\phi"] & \mathcal{B}(\overline{V})\#k[\overline{\Gamma}] \arrow[r] & 1 \end{tikzcd}$$

where ϕ is defined by $\phi(hx_1^{k_1}x_2^{k_2}\dots x_t^{k_t}) = \overline{h}\overline{v_1}^{k_1}\overline{v_2}^{k_2}\dots\overline{v_t}^{k_t}$. By Corollary 2.3 the dual of $\mathcal{B}(\overline{V})\#k[\overline{\Gamma}]$ is $\mathcal{B}(W)\#k[\Omega]$ where $W = \oplus_{i=1}^{t} kw_i$ with $w_i \in W_{\chi_i}^{\overline{g_i}}$ and so $\mathcal{B}(W)\#k[\Omega]$ is isomorphic to a pointed sub Hopf algebra of A^*. We denote this subHopf algebra by B and note that $C(\overline{1}) = \mathcal{B}(W)\#k[\Theta]$ is a subHopf algebra of B.

(3) Since $k[\Theta] \subseteq C(\overline{1})$, a subHopf algebra of A^*, $C(\overline{\gamma})$ is well-defined and Lemma 3.1 implies $\gamma C(\overline{1}) = C(\overline{1})\gamma$. If γ is grouplike, then the map $z \to \gamma z$ from $C(\overline{1})$ to $\gamma C(\overline{1})$ is a coalgebra isomorphism. If $\gamma \notin \Omega$, then $\triangle(\gamma) = \gamma\otimes\gamma + z(\gamma) \in A^*\otimes A^*$. Here $z(\gamma)$ is a sum of terms of the form $\xi w(\lambda, x_1^{n_1}\dots x_t^{n_t}) \otimes w(\lambda', x_1^{m_1}\dots x_t^{m_t})$ where we have $\xi\lambda(h)\lambda'(l) = \gamma(hx_1^{n_1}\dots x_t^{n_t} l x_1^{m_1}\dots x_t^{m_t}) = \lambda''(l)\gamma(hlx_1^{n_1}\dots x_t^{n_t}x_1^{m_1}\dots x_t^{m_t})$ where $\lambda'' \in \Theta$. Since this holds for all h, l we must have $\xi = \gamma(x_1^{n_1}\dots x_t^{n_t}x_1^{m_1}\dots x_t^{m_t})$ and $\lambda = \gamma$, $\lambda' = \gamma\lambda''$ with $\lambda'' \in \Theta$. Thus $w(\lambda, x_1^{n_1}\dots x_t^{n_t}) \otimes w(\lambda', x_1^{m_1}\dots x_t^{m_t}) \in C(\overline{1})\gamma \otimes C(\overline{1})\gamma$ and $C(\overline{\gamma})$ is a coalgebra as required. Clearly $C(\overline{1})\gamma$ has k basis $\mathcal{W}_\gamma = \{w(\gamma\chi, z) | \chi \in \Theta, z \in \mathcal{Z}\}$ so that $A^* = \oplus_{\overline{\gamma}\in\hat{\Gamma}/\Theta} C(\overline{\gamma})$ as required.

(4) Since $C(\overline{1})$ is a subHopf algebra,

$$C(\overline{\gamma})C(\overline{\lambda}) = \gamma C(\overline{1})C(\overline{1})\lambda = \gamma C(\overline{1})\lambda = (\gamma\lambda)C(\overline{1}) = C(\overline{\gamma\lambda}).$$

If $\gamma \in \hat{\Gamma}$, since $S_{A^*}(\gamma) = (S_A)^*(\gamma) = \gamma^{-1}$, the last statement is clear.

□

We describe A^* when $\dim V = 1$. It is shown in [31, 2.3] using the algebra structure of A that for $\Gamma =< g >$ cyclic of order np and $V = kv = V_g^{\chi^p}$ where χ generates $\hat{\Gamma}$, then for A the unique nontrivial lifting of $\mathcal{B}(V)\#k[\Gamma]$, the dual A^* contains $p-1$ matrix coalgebras of dimension n^2. Also it is shown in [17] that A^* contains a nontrivial matrix subcoalgebra. We generalize this description to Γ any finite abelian group and explicitly calculate the matrix coalgebras.

Theorem 3.3. *Let* $V = kv \in {}^{\Gamma}_{\Gamma}\mathcal{YD}$ *with* $v \in V_g^{\chi}$ *and* $\chi(g)$ *a primitive* r*th root of unity. Suppose* $\chi^r = \epsilon, g^r \neq 1$ *and* A *is the (only) nontrivial lifting of* $\mathcal{B}(V)\#k[\Gamma]$*. Then as a coalgebra,* $A^* \cong \oplus_{\overline{\gamma}\in\hat{\Gamma}/\Theta} C(\overline{\gamma})$ *where* $C(\overline{\gamma}) \cong \mathcal{M}^c(r,k)$ *if* $\gamma \notin \Omega$ *and* $C(\overline{\gamma}) \cong C(\overline{1}) \cong \mathcal{B}(W)\#k[\Theta]$ *as coalgebras with* $W = k\cdot w, w \in W_{\chi}^{\overline{g}}$ *if* $\gamma \in \Omega$*. Thus corad* (A^*) *is the sum of the group algebra* $k[\Omega]$ *and* m $r\times r$ *matrix coalgebras where* $mr = |\hat{\Gamma}| - |\Omega|$*. As an algebra* A^* *is generated by* $\hat{\Gamma}$ *and* w *where* $\gamma * w = \gamma(g)w * \gamma$ *for* $\gamma \in \hat{\Gamma}$ *and* $w^r = 0$*.*

Proof. Let $\gamma \in \hat{\Gamma} - \Omega$ and we show that $C(\overline{\gamma})$ is an $r\times r$ matrix coalgebra by explicitly exhibiting a matrix coalgebra basis. For $1 \leq i, j \leq r$, define $e(i,j) = w(\gamma\chi^{i-1}, x^{i-j})$ if $i \geq j$ and $e(i,j) = \eta w(\gamma\chi^{i-1}, x^{r+i-j})$ if $i < j$ where $\eta = \gamma(g^r - 1) \neq 0$. Then we claim that $\triangle(e(j,k)) = (m_A^*)(e(j,k))$ is $\sum_{s=1}^{r} e(j,s) \otimes e(s,k)$. First we note that for $j \geq k$,

$$\begin{aligned}
\triangle w(\gamma\chi^{j-1}, x^{j-k}) &= \sum_{s=1}^{k-1} w(\gamma\chi^{j-1}, x^{j-s}) \otimes \eta w(\gamma\chi^{s-1}, x^{r+s-k}) \\
&+ \sum_{s=k}^{j} w(\gamma\chi^{j-1}, x^{j-s}) \otimes w(\gamma\chi^{s-1}, x^{s-k}) \\
&+ \sum_{s=j+1}^{r} \eta w(\gamma\chi^{j-1}, x^{r+j-s}) \otimes w(\gamma\chi^{s-1}, x^{s-k}).
\end{aligned}$$

Now in the first sum, $s < k \leq j$ and so we have $e(j,s) = w(\gamma\chi^{j-1}, x^{j-s})$ and $e(s,k) = \eta w(\gamma\chi^{s-1}, x^{r+s-k})$, and the sum is $\sum_{s=1}^{k-1} e(j,s) \otimes e(s,k)$. The others are similar and we conclude that $\triangle(w(\gamma\chi^{j-1}, x^{j-k})) = \triangle(e(j,k)) = \sum_{s=1}^{r} e(j,s) \otimes e(s,k)$.

Similarly, if $j < k$,

$$\begin{aligned}
\triangle \eta w(\gamma\chi^{j-1}, x^{r+j-k}) &= \eta[\sum_{s=1}^{j} w(\gamma\chi^{j-1}, x^{j-s}) \otimes w(\gamma\chi^{s-1}, x^{r+s-k})] \\
&+ \eta[\eta \sum_{s=j+1}^{k-1} w(\gamma\chi^{j-1}, x^{j-s}) \otimes w(\gamma\chi^{s-1}, x^{r+s-k})] \\
&+ \eta[\sum_{s=k}^{r} (\gamma\chi^{j-1}, x^{r+j-s}) \otimes w(\gamma\chi^{s-1}, x^{s-k})].
\end{aligned}$$

Note that the middle sum is empty if $j+1 = k$. It is easily verified that this sum is $\sum_{s=1}^{r} e(j,s) \otimes e(s,k)$. The last statement follows from Lemma 3.1. □

Remark 3.4. *For A, V, Γ, g, χ as in Theorem 3.3, Jac(A) is the ideal I generated by $(1 + g^r + \ldots + g^{(s-1)r})x$ where s is the order of g^r. For I is clearly contained in Jac(A) since I is a nil ideal and* $\dim I = |\Gamma / < g^r > |(r-1) = |\Omega|(r-1)$, *while* $\dim \mathit{corad}(A^*) = |\Omega| + r(|\Gamma| - |\Omega|) = \dim A - |\Omega|(r-1)$. □

Corollary 3.5. *Let A be a lifting of $\mathcal{B}(V)\#k[\Gamma]$ where $V = \oplus_{i=1}^{t} kv_i$ is a quantum linear space and x_i is the lifting of $v_i\#1$ to A. Suppose that the matrix $\mathcal{A}$ has $\alpha_{1,1} = 1$ and has all other entries 0, i.e. $x_i x_j = \chi_j(g_i) x_j x_i$, for $1 \leq i, j \leq t$, and $x_j^{r_j} = 0$ for $j = 2, \ldots, t$ while $x_1^{r_1} = g_1^{r_1} - 1$. Then A^* has coradical $C \cong k[\Omega] \oplus \oplus_{\substack{\overline{\gamma} \in \hat{\Gamma}/\Theta \\ \gamma \notin \Omega}} \mathcal{M}^c(r_1, k)$.*

Proof. Let I be the Hopf ideal of A generated by the skew-primitives $x_2, \ldots, x_t$. Then there is a Hopf algebra map from A onto $A_1 = A/I$, and A_1 is isomorphic to the lifting of $\mathcal{B}(kv_1)\#k[\Gamma]$ considered in Theorem 3.3. Thus A^* has a sub Hopf algebra isomorphic to A_1^* and by Theorem 3.3, the coradical of A_1^* is isomorphic to C. Now, Jac(A) contains the nil ideals $N_1 =< x_2, \ldots, x_t >$ and $N_2 =< (1 + g_1^{r_1} + \ldots + g_1^{(m_1-1)r_1})x_1 >$ where $m_1 r_1$ is the order of g_1. Thus $\dim Jac(A) \geq |\Gamma| r_1 (r_2 \ldots r_t - 1) + |\Omega|(r_1 - 1) r_2 \ldots r_t - |\Omega|(r_1 - 1)(r_2 \ldots r_t - 1) = |\Gamma| r_1 (r_2 \ldots r_t - 1) + |\Omega|(r_1 - 1)$, i.e. Jac($A$) has dimension greater than or equal to $\dim N_1 + \dim N_2 -$

$\dim(N_1 \cap N_2)$. Thus $\dim(coradA^*) \leq |\Gamma| r_1 r_2 \ldots r_t - |\Gamma| r_1(r_2 \ldots r_t - 1) - |\Omega|(r_1 - 1)$ $= |\Gamma| r_1 - |\Omega|(r_1 - 1) = |\Omega| + (|\Gamma| - |\Omega|) r_1 = \dim C$. □

Now we consider quantum linear spaces of dimension 2. The next result is proved in [31] for the group Γ cyclic but without our restrictions on r. The approach is rather different.

Theorem 3.6. *Let $V = kv_1 \oplus kv_2$ be a quantum linear space with $v_1 \in V_g^{\chi}$ and $v_2 \in V_{g^n}^{\chi^m}$ and such that*

i) $\chi(g)$ and $\chi^m(g^n)$ are primitive rth roots of unity for r an odd squarefree integer.

ii) χ and χ^m have order r in the group $\hat{\Gamma}$, i.e. $(m, r) = 1 = (n, r)$.

iii) $g^r \neq 1$ and $g^{nr} \neq 1$.

Let A be the lifting of $\mathcal{B}(V)\#k[\Gamma]$ with matrix $\mathcal{A} = I_2$. Then for $\gamma \notin \Omega = G(A^*)$, $C(\overline{\gamma}) \cong \oplus_{i=1}^r \mathcal{M}^c(r, k)$, and $corad(A) \cong k[\Omega] \oplus \oplus_{\substack{\overline{\gamma} \in \hat{\Gamma}/\Theta \\ \gamma \notin \Omega}} (\oplus_{i=1}^r \mathcal{M}^c(r, k))$.

Proof. Let x and y be the liftings of v_1 and v_2 to A. Then $yx = q^n xy$ where $q = \chi(g)$ is a primitive rth root of unity and $x^r = g^r - 1$, $y^r = g^{nr} - 1$. The dimension of $C(\overline{\gamma})$ for any $\overline{\gamma}$ is r^3.

Let $\gamma \notin \Omega = G(A^*)$. Since $C(\overline{\gamma})$ has no grouplike elements, $C(\overline{\gamma})$ must contain a matrix coalgebra $M \cong \mathcal{M}^c(s, k)$ for some integer $s \geq 2$. Thus M has a matrix coalgebra basis $e(i,j), 1 \leq i, j \leq s$ with $\triangle(e(i,i)) = e(i,i) \otimes e(i,i) + z_i \in C(\overline{\gamma}) \otimes C(\overline{\gamma})$ and $\epsilon(e(i,i)) = 1$.

The only elements z in the basis $\mathcal{W}_\gamma = \{w(\gamma\chi^i, x^j y^k),\ 0 \leq i, j, k \leq r-1\}$ for $C(\overline{\gamma})$, such that a scalar multiple of $z \otimes z$ is a summand of $\triangle w$ for some basis element $w \in \mathcal{W}_\gamma$ are those of the form $z = w(\gamma\chi^i, x^j y^k)$ where $\chi^j \chi^{mk} = \epsilon$, i.e., $x^j y^k$ commutes with elements of Γ. Thus for every l, $1 \leq l \leq s$, then $e(l,l)$ is a linear combination of such elements. If $e(l,l) = \sum_{i=0}^{r-1} a_i \gamma\chi^i + w$, for w a linear combination of basis elements from $\mathcal{W}_\gamma \setminus \hat{\Gamma}$, then it is clear that exactly one a_i is 1 and the rest are 0 since terms of the form $\gamma\chi^i \otimes \gamma\chi^j$ with $i \neq j$ do not occur in $\triangle(z)$ for any $z \in A^*$. For $w = w(\gamma\chi^i, x^j y^k) \in \mathcal{W}_\gamma$, $\triangle w = \gamma\chi^i \otimes w + w \otimes \gamma\chi^i\chi^{-j}\chi^{-km} + w'$ where w' is a linear combination of terms from $\mathcal{W}_\gamma \setminus \hat{\Gamma} \otimes \mathcal{W}_\gamma \setminus \hat{\Gamma}$. Thus $e(l,l) = \lambda + \sum_{i=1}^{r-1} a_i w(\lambda, x^{nr-im} y^i) = \lambda + \sum_{i=1}^{r-1} a_i z_{i,\lambda}$, for some $\lambda \in \overline{\gamma}$, where $z_{i,\lambda} = w(\lambda, x^{nr-im} y^i)$ with n the integer such that $0 \leq nr - im \leq r-1$. (This also shows that the $e(i,j), i \neq j$, are linear combinations of elements of $\mathcal{W}_\gamma \setminus \hat{\Gamma}$.) Now, in the expression for $\triangle(\lambda)$, a nonzero scalar multiple of $w(\lambda, x^j) \otimes w(\lambda\chi^{-j}, x^{r-j})$ occurs for all $1 \leq j \leq r-1$. Also no such summand occurs in $\triangle z_{i,\lambda}$ for any i and so $w(\lambda, x^j)$ must be a summand in some $e(l,m)$. But then $e(m,m)$ must contain $\lambda\chi^{-j}$. Then, for all $\theta \in \overline{\gamma}$, θ is a summand of some matrix coalgebra basis element $e(m,m)$. Thus the dimension of M is at least r^2.

Since χ^i is grouplike, the map $z \to \chi^i z \chi^{-i}$ from $C(\overline{\gamma})$ to $C(\overline{\gamma})$ is a coalgebra isomorphism. Suppose the r subcoalgebras $\chi^i M \chi^{-i}$ of $C(\overline{\gamma})$, $i = 0, 1, \ldots, r-1$, are not all distinct (this will be the case if $s > r$) and so $\chi^i M \chi^{-i} = M$ for some i.

Let $\Theta' = \{\chi^i | \chi^i M \chi^{-i} = M\} \subseteq <\chi>$, and let k to be the smallest positive integer such that $\chi^k \in \Theta'$. Then $\Theta' = <\chi^k>$ and $r = kt$ for some t.

For $\theta, \lambda \in C(\overline{\gamma})$, denote by $E(\theta, \lambda)$ the subspace of M spanned by those $e(i,j)$ such that $\triangle e(i,j)$ contains $\theta \otimes e(i,j)$ and $e(i,j) \otimes \lambda$ as summands. By the previous discussion, $\dim E(\theta, \lambda) \geq 1$ for all $\theta, \lambda \in \overline{\gamma}$. Note that $\dim E(\theta, \theta)$ is a square, since if $e(i,i), e(j,j) \in E(\theta, \theta)$ then $e(i,j), e(j,i) \in E(\theta, \theta)$. Conversely, if $e(i,j) \in E(\theta, \theta)$, then $e(i,i)$ and $e(j,j)$ lie in $E(\theta, \theta)$. Also if $\dim E(\theta, \theta) = l^2$ and $\dim E(\lambda, \lambda) = m^2$ then $\dim E(\theta, \lambda) = \dim E(\lambda, \theta) = lm$. This is because if $e(i,i) \in E(\theta, \theta)$ and $e(j,j) \in E(\lambda, \lambda)$ then $e(i,j) \in E(\theta, \lambda)$ and $e(j,i) \in E(\lambda, \theta)$ and conversely. Note that $\chi^k E(\theta, \lambda) \chi^{-k} = E(\theta, \lambda)$ for all $\theta, \lambda \in \overline{\gamma}$.

Let $E(\theta, \theta)$ contain a vector v such that $v = a_0\theta + \sum_{i=1}^{r-1} a_i z_{i,\theta}$ for $z_{i,\theta} = w(\theta, x^{nr-im}y^i)$ as above, with $a_i \neq 0$ for $0 \leq i \leq r-1$. Now $\chi^k v \chi^{-k} = a_0\theta + \sum_{i=1}^{r-1} \chi^k(g^{nr-im}g^{in}) a_i z_i$ where $\chi^k(g^{nr-im}g^{in}) = q^{ki(m-n)}$. Since V is a quantum linear space, $q^{m+n} = 1$ and so $m + n = pr$ for some p and $m - n = 2m - pr$. Thus $q^{ki(m-n)} = (q^{2m})^{ki}$ and since $(2, r) = (m, r) = 1$, q^{2m} is a primitive rth root of unity, and $q' = q^{2mk}$ is a primitive tth root of unity. Thus the t vectors $\chi^{ik} v \chi^{-ik}$, $0 \leq i \leq t-1$, are linearly independent since the rank of the Vandermonde matrix $V(1, q', q'^2, \ldots, q'^{t-1})$ is t. So $\dim E(\theta, \theta) \geq t$, and since if $t > 1, t$ is not a square by the assumption that r is square-free, $\dim E(\theta, \theta) > t$. Thus, if all $E(\theta, \theta)$ contain such a vector v, then $\dim M = \sum_{\theta, \lambda \in \overline{\gamma}} \dim E(\theta, \lambda) > r^2 t$. Since the isomorphic coalgebras $M, \chi M \chi^{-1}, \ldots, \chi^{k-1} M \chi^{-(k-1)}$ are distinct by our choice of k, $\dim C(\overline{\gamma}) > r^2 tk = r^3$, which is a contradiction. Thus $k = r, t = 1$, M has dimension r^2 and $C(\overline{\gamma}) \cong \oplus_{j=1}^r \mathcal{M}^c(r, k)$.

It remains to show that every $E(\theta, \theta)$ contains a vector v of the required form.

Let $e(l,l) \in E(\theta, \theta)$ with $\dim E(\theta, \theta) > 1$ and suppose $a_1 = 0$ in the expression for $e(l,l)$. Since $\triangle\theta$ is a summand of $\triangle e(l,l), \zeta z_{1,\theta} \otimes z_{r-1,\theta}$ is a summand of $\triangle e(l,l)$, and $\zeta' z_{1,\theta}$ is a summand of $e(l,j)$ for some $j \neq l$, with $e(l,j), e(j,j), e(j,l) \in E(\theta, \theta)$. (Here ζ, ζ', etc. denote nonzero scalars.) Similarly, a nonzero multiple of $z_{2,\theta}$ is a summand in $e(l,l)$ or in some $e(l,j) \in E(\theta, \theta)$. Continuing in this way we find a vector v in $E(\theta, \theta)$ as required, by taking some linear combination of $e(l,l)$ and the $e(l,j)$. □

Remark 3.7. *In the setting of Theorem 3.6, if* $r = 2$, *then* $q^{2nki} = 1$ *and* $\chi^k e(\lambda, \lambda) \chi^{-k} = e(\lambda, \lambda)$ *so that* M *is invariant under the action of* χ. □

Corollary 3.8. *For* A *as in Theorem 3.6,* $I = \{h(1 + g^r + \ldots + g^{r(l-1)})x^i y^j | h \in \Gamma$, l *is the order of* g^r *in* Γ, $0 \leq i, j \leq r-1$ *and* $i + j > 0\}$ *is the Jacobson radical of* A.

Proof. Clearly I is a nil ideal in A and has dimension $|\Omega|(r^2 - 1)$. But

$$\dim Jac(A) = |\Gamma| r^2 - \dim corad(A^*) = |\Gamma| r^2 - (|\Omega| + \left(\frac{|\Gamma| - |\Omega|}{r}\right) r^3) = |\Omega|(r^2 - 1).$$

□

Corollary 3.9. *Let* A *be a lifting of* $\mathcal{B}(V)\#k[\Gamma]$ *where* $V = \oplus_{i=1}^t kv_i$ *is a quantum linear space such that* $V' = kv_1 \oplus kv_2$ *satisfies the conditions in Theorem 3.6, and*

A has lifting matrix $\mathcal{A} = \left[\begin{array}{c|c} I_2 & 0 \\ \hline 0 & 0 \end{array}\right]$. Then in $A^*, coradC(\overline{\gamma}) \cong \oplus_{i=1}^{r}\mathcal{M}^c(r,k)$ for $\gamma \notin \Omega$.

Proof. Let $N_1 = \{h(1+g^r+\ldots+g^{r(l-1)})x_1^{m_1}x_2^{m_2}\ldots x_t^{m_t} | h \in \Gamma$, l is the order of g^r, $0 \leq m_i \leq r_i - 1$ and $m_1 + m_2 > 0\}$ and let $N_2 = \{hx_1^{m_1}x_2^{m_2}x_3^{m_3}\ldots x_t^{m_t} | h \in \Gamma, 0 \leq m_i \leq r_i - 1, m_3 + \ldots + m_t > 0\}$. Since N_1 and N_2 are nil ideals of A, then $Jac(A)$ has dimension at least $|\Omega|(r^2-1)r_3\ldots r_t + |\Gamma|r^2(r_3\ldots r_t - 1) - |\Omega|(r^2-1)(r_3\ldots r_t - 1) = |\Gamma|r^2(r_3\ldots r_t - 1) + |\Omega|(r^2-1)$. Thus $\dim corad(A^*) \leq |\Gamma|r^2 r_3\ldots r_t - |\Gamma|r^2(r_3\ldots r_t - 1) - |\Omega|(r^2-1) = (|\Gamma| - |\Omega|)r^2 + |\Omega|$. Now, as in the proof of Corollary 3.5, let I be the Hopf ideal of A generated by the skew-primitives $x_3, \ldots, x_t$. Then $A_1 = A/I$ is isomorphic to the lifting of $\mathcal{B}(kv_1 \oplus kv_2)\#k[\Gamma]$ with matrix $\mathcal{A} = I_2$ and so $A_1^* \subseteq A^*$ has coradical isomorphic to $k[\Omega] \oplus \oplus_{\substack{\overline{\gamma} \in \hat{\Gamma}/\Theta \\ \gamma \notin \Omega}} (\oplus_{i=1}^{r}\mathcal{M}^c(r,k))$ by Theorem 3.6. But the dimension of this space is $|\Omega| + (\frac{|\Gamma| - |\Omega|}{r})r^3 = |\Omega| + (|\Gamma| - |\Omega|)r^2$. □

In the next section we do some detailed computations of bases for matrix coalgebras in the duals of some liftings of quantum linear spaces.

4. Examples

4.1. **Duals with grouplikes of order 3.** The two examples in this subsection illustrate Theorem 3.6, i.e. they are duals of liftings of quantum linear spaces with matrix $\mathcal{A} = I_2$.

Remark 4.1. In the setting of Theorem 3.6, in order to choose $e(i,i)$ in a matrix coalgebra basis, each summand αw of $e(i,i)$ must be one such that $\alpha w \otimes \alpha w$ can occur as a summand in $\triangle(\beta w')$ where w, w' lie in a basis $\mathcal{W}$ for A^*. This was noted in the proof of Theorem 3.6.

Suppose $v_1 \in V_g^{\chi}$ and $v_2 \in V_g^{\chi^{-1}}$ so that in $C(\overline{\gamma})$, $e(1,1) = \sum_{i=0}^{r-1} a_i w(\gamma, (xy)^i)$. Then $a_0 = 1$. Clearly $a_2 = a_1^2, a_4 = a_2^2$, etc. If $r = 3$, then $a_2 = a_1^2$ and $\eta^2 a_1 = a_2^2$ where $\eta = \gamma(g^3 - 1)$. Then $\eta^2 a_1 = a_1^4$ so $a_1^3 = \eta^2$ and $a_1 = q\eta^{2/3}$ where $q^3 = 1$. Similar computations can be done for other values of r. For instance, if $r = 5$, then $a_2 = a_1^2, a_4 = a_2^2, \eta^2 a_3 = a_4^2, \eta^2 a_1 = a_3^2$ where here $\eta = \gamma(g^5 - 1)$. Then $\eta^6 a_1 = \eta^4 a_3^2 = a_4^4 = a_2^8 = a_1^{16}$ and thus $a_1^{15} = \eta^6$ so if $a_1 = q\eta^{6/15} = a\eta^{2/5}$ where $q^5 = 1$ then $a_i = q^i \eta^{2i/5}$. □

Example 4.2. For Γ finite abelian as usual, let $V = kv_1 \oplus kv_2$ with $v_1 \in V_g^{\chi}, v_2 \in V_g^{\chi^{-1}}$ where $\chi^3 = \epsilon$ and $g^3 \neq 1$. Let A be the nontrivial lifting of $\mathcal{B}(V)\#k[\Gamma]$ with matrix $\mathcal{A} = I_2$. Denote by x, y the liftings of the $v_i\#1$ to A and write $q = \chi(g)$, a primitive cube root of unity. Then $yx = qxy$, $x^3 = y^3 = g^3 - 1$. By Proposition 3.2, A^* contains a subHopf algebra of dimension 27 isomorphic to $\mathcal{B}(W)\#k[\Theta]$ and this subHopf algebra is denoted $C(\overline{1})$. We compute the matrix coalgebras that constitute $\gamma C(\overline{1})$ for $\gamma \notin \Omega$. We work with the k-basis $w(\lambda, z)$ where $\lambda \in \hat{\Gamma}$ and $z \in \mathcal{Z} = \{1, x, y, x^2, y^2, xy, x^2y, xy^2, (xy)^2\}$.

Then a k-basis for $C(\overline{\gamma})$ is $w(\gamma\chi^i, z)$ for $0 \leq i \leq 2$ and $z \in \mathcal{Z}$ as above. As in the proof of Theorem 3.6, since the only elements of $\mathcal{Z}$ that commute with elements of Γ are $1, xy$ and $(xy)^2$, then any candidate for a matrix coalgebra basis element $e(i,i)$ containing γ as a summand must be a linear combination of $\gamma, w(\gamma, xy)$ and $w(\gamma,(xy)^2)$. Let $\eta = \gamma(g^3-1) \neq 0$. Supplying extra detail in this first example, we note that

$$\begin{array}{rl} & \triangle(\gamma) = \gamma \otimes \gamma + \eta[w(\gamma,x) \otimes w(\gamma\chi^2,x^2) + w(\gamma,x^2) \otimes w(\gamma\chi,x) \\ + & w(\gamma,y) \otimes w(\gamma\chi,y^2) + w(\gamma,y^2) \otimes w(\gamma\chi^2,y)] + \eta^2[qw(\gamma,xy^2) \otimes w(\gamma\chi,x^2y) \\ + & qw(\gamma,x^2y) \otimes w(\gamma\chi^2,xy^2) + w(\gamma,xy) \otimes w(\gamma,(xy)^2) + w(\gamma,(xy)^2) \otimes w(\gamma,xy)]; \end{array}$$

$$\begin{array}{rl} & \triangle(w(\gamma,xy)) = \gamma \otimes w(\gamma,xy) + w(\gamma,x) \otimes w(\gamma\chi^2,y) + qw(\gamma,y) \otimes w(\gamma\chi,x) \\ + & \eta[w(\gamma,x^2) \otimes w(\gamma\chi,x^2y) + q^2w(\gamma,y^2) \otimes w(\gamma\chi^2,xy^2) + w(\gamma,xy^2) \otimes w(\gamma\chi,y^2) \\ + & q^2w(\gamma,x^2y) \otimes w(\gamma\chi^2,x^2)] + \eta^2 w(\gamma,(xy)^2) \otimes w(\gamma,(xy)^2) + w(\gamma,xy) \otimes \gamma; \end{array}$$

$$\begin{array}{rl} & \triangle(w(\gamma,(xy)^2) = \gamma \otimes w(\gamma,(xy)^2) + w(\gamma,xy) \otimes w(\gamma,xy) + qw(\gamma,y) \otimes w(\gamma\chi,x^2y) \\ + & qw(\gamma,xy^2) \otimes w(\gamma\chi,x) + q^2w(\gamma,x^2) \otimes w(\gamma\chi,y^2) + q^2w(\gamma,x) \otimes w(\gamma\chi^2,xy^2) \\ + & w(\gamma,y^2) \otimes w(\gamma\chi^2,x^2) + q^2w(\gamma,x^2y) \otimes w(\gamma\chi^2,y) + w(\gamma,(xy)^2) \otimes \gamma. \end{array}$$

Now define $e(1,1) = \gamma + \eta^{2/3}w(\gamma,xy) + \eta^{4/3}w(\gamma,(xy)^2)$ as suggested by Remark 4.1. By the computations above, $\triangle(e(1,1) = e(1,1) \otimes e(1,1) + e(1,2) \otimes e(2,1) + e(1,3) \otimes e(3,1)$, where

$$\begin{array}{rcl} e(1,2) & = & \eta^{1/3}w(\gamma,x) + \eta^{2/3}w(\gamma,y^2) + q^2\eta w(\gamma,x^2y); \\ e(2,1) & = & \eta^{2/3}w(\gamma\chi^2,x^2) + \eta^{1/3}w(\gamma\chi^2,y) + q^2\eta w(\gamma\chi^2,xy^2); \\ e(1,3) & = & \chi^2 e(2,1)\chi^2 = q\eta^{2/3}w(\gamma,x^2) + q^2\eta^{1/3}w(\gamma,y) + q^2\eta w(\gamma,xy^2); \\ e(3,1) & = & \chi^2 e(1,2)\chi^2 = q^2\eta^{1/3}w(\gamma\chi,x) + q\eta^{2/3}w(\gamma\chi,y^2) + q^2\eta w(\gamma\chi,x^2y). \end{array}$$

Now we compute $\triangle(e(3,1))$ and define $e(3,3) = \gamma\chi + q\eta^{2/3}w(\gamma\chi,xy) + q^2\eta^{4/3}w(\gamma\chi,(xy)^2) = \chi^2 e(1,1)\chi^2$. Similarly we have

$$\begin{array}{rcl} e(2,2) & = & \chi e(1,1)\chi = \gamma\chi^2 + q^2\eta^{2/3}w(\gamma\chi^2,xy) + q\eta^{4/3}w(\gamma\chi^2,(xy)^2); \\ e(2,3) & = & \chi e(1,2)\chi = q\eta^{1/3}w(\gamma\chi^2,x) + q^2\eta^{2/3}w(\gamma\chi^2,y^2) + q^2\eta w(\gamma\chi^2,x^2y); \\ e(3,2) & = & \chi e(2,1)\chi = q^2\eta^{2/3}w(\gamma\chi,x^2) + q\eta^{1/3}w(\gamma\chi,y) + q^2\eta w(\gamma\chi,xy^2). \end{array}$$

Straightforward computation then shows that these elements satisfy the conditions to be a basis for a 3×3 matrix coalgebra and that by the proof of Theorem 3.6 or by easy computation, this matrix coalgebra is not invariant under the inner action of χ. Thus the other two matrix coalgebras constituting $C(\overline{\gamma})$ may be obtained by using this action and $C(\overline{\gamma}) = \oplus_{i=1}^{3}\mathcal{M}^c(3,k)$. □

Example 4.3. This example is similar to Example 4.2 but $m = 1$ and $n = -1$, i.e. $V = kv_1 \oplus kv_2$ with $v_1 \in V_g^\chi$ and $v_2 \in V_{g^{-1}}^\chi$ where $\chi^3 = \epsilon$ and $g^3 \neq 1$. Otherwise $A, \mathcal{A}, x, y, q$ are as defined in Example 4.2. Again we compute the matrix coalgebras that constitute $C(\overline{\gamma})$ for $\gamma \notin \Omega = G(A^*)$. This time, we work with the k-basis for A given by hz, $h \in \Gamma$, $z \in \mathcal{Z} = \{1, x, y, x^2, y^2, xy, x^2y, xy^2, x^2y^2\}$ so A^* has basis $w(\gamma\chi^k, x^iy^j), 0 \leq k, i, j \leq 2$. Here the elements of $\mathcal{Z}$ that commute with grouplikes are h, hxy^2 or $hx^2y, h \in \Gamma$, so that we guess that $e(1,1)$ has the form

$\gamma + aw(\gamma, x^2y) + bw(\gamma, xy^2)$. Straightforward computation yields that $a^3 = b^3 = (\eta q)^3 = \eta^3$. We choose $a = b = \eta q$, and then find that

$$\begin{aligned}
e(1,1) &= \gamma + \eta q w(\gamma, x^2y) + \eta q w(\gamma, xy^2);\\
e(2,2) &= \gamma\chi^2 + \eta q^2 w(\gamma\chi^2, x^2y) + \eta w(\gamma\chi^2, xy^2) = \chi e(1,1)\chi;\\
e(3,3) &= \gamma\chi + \eta w(\gamma\chi, x^2y) + \eta q^2 w(\gamma\chi, xy^2) = \chi^2 e(1,1)\chi^2;\\
e(1,2) &= \eta w(\gamma, x) + \eta q^2 w(\gamma, y) + \eta^2 w(\gamma, x^2y^2);\\
e(3,1) &= \eta q^2 w(\gamma\chi, x) + \eta w(\gamma\chi, y) + \eta^2 w(\gamma\chi, x^2y^2) = \chi^2 e(1,2)\chi^2;\\
e(2,3) &= \eta^{-1} q w(\gamma\chi^2, x) + \eta^{-1} q w(\gamma\chi^2, y) + w(\gamma\chi^2, x^2y^2) = \eta^{-2}\chi e(1,2)\chi;\\
e(2,1) &= w(\gamma\chi^2, x^2) + q w(\gamma\chi^2, y^2) + q w(\gamma\chi^2, xy);\\
e(1,3) &= q w(\gamma, x^2) + w(\gamma, y^2) + q w(\gamma, xy) = \chi^2 e(2,1)\chi^2;\\
e(3,2) &= q^2\eta^2 w(\gamma\chi, x^2) + \eta^2 q^2 w(\gamma\chi, y^2) + \eta^2 q w(\gamma\chi, xy) = \eta^2 \chi e(2,1)\chi;
\end{aligned}$$

form a matrix coalgebra basis for a matrix coalgebra $E = \chi E \chi = \chi^2 E \chi^2$. Then $C(\overline{\gamma}) = E \oplus E\chi \oplus E\chi^2 \cong \oplus_{i=1}^{3} \mathcal{M}^c(3,k)$. □

4.2. **Duals of pointed Hopf algebras of dimension 16.** Now we describe the duals of all pointed Hopf algebras of dimension 16 which are not group algebras or their duals. In fact, this means that the duals of all Hopf algebras of dimension 16 with coradical a Hopf algebra are known, since the nonpointed Hopf algebras of dimension 16 with coradical a proper subHopf algebra were computed in [12] and found to be self-dual. In [11], all pointed Hopf algebras of dimension 16 over an algebraically closed field k of characteristic 0 were computed, up to isomorphism. The classification of semisimple Hopf algebras of dimension 16 is done in [20].

By [11], there are 43 different isomorphism classes of pointed Hopf algebras: 14 for the group algebras and 29 for Hopf algebras with group of grouplikes of order 2, 4 or 8. Of these 29 Hopf algebras, 21 are of the form $\mathcal{B}(V)\#k[\Gamma]$ where V is a quantum linear space and 8 are nontrivial liftings of such $\mathcal{B}(V)\#k[\Gamma]$. First we list the Radford biproducts and their duals. We use the following notation.

If Γ is cyclic, then its generator is denoted by c and $\hat{\Gamma}$ has generator c^*. If Γ is the product of 2 cyclic groups then we denote their generators by c of order n and d of order m where $n \geq m$ and c^*, d^* are the elements of $\hat{\Gamma}$ which map c to a primitive nth root of unity and d to 1, d to a primitive mth root of unity and c to 1, respectively. If Γ is the product of 3 cyclic groups, we denote the generators by c, d, e and the generators of $\hat{\Gamma}$ by c^*, d^*, e^* with the same conventions.

We describe the space $V = \oplus_{j=1}^{\dim V} kv_j$ with $v_j \in V_{g_j}^{\chi_j}$ by pairs (g_j, χ_j), one pair for each v_j. A list of the $H_i = \mathcal{B}(V_i)\#k[\Gamma_i]$ and their duals is given by the table below.

H_i	Γ_i	V_i	Dual of H_i
H_1	C_8	(c, c^{*4})	H_2
H_2	C_8	(c^4, c^*)	H_1
H_3	C_8	(c^2, c^{*2})	self-dual
H_4	C_8	$(c^6, c^{*2}) \cong (c^2, c^{*6})$	self-dual
H_5	$C_4 \times C_2$	(c, c^{*2})	H_6
H_6	$C_4 \times C_2$	(c^2, c^*)	H_5
H_7	$C_4 \times C_2$	(cd, d^*)	H_8
H_8	$C_4 \times C_2$	$(d, c^*d^*) \cong (c^2d, c^*)$	H_7
H_9	$C_4 \times C_2$	(d, d^*)	self-dual
H_{10}	$C_2 \times C_2 \times C_2$	(c, c^*)	self-dual
H_{11}	C_4	(c, c^*)	self-dual
H_{12}	C_4	$(c, c^{*3}) \cong (c^3, c^*)$	self-dual
H_{13}	C_4	$(c^2, c^*)(c^2, c^*)$	H_{14}
H_{14}	C_4	$(c, c^{*2})(c, c^{*2})$	H_{13}
H_{15}	C_4	$(c^2, c^*)(c^2, c^{*3})$	H_{16}
H_{16}	C_4	$(c, c^{*2})(c^3, c^{*2})$	H_{15}
H_{17}	$C_2 \times C_2$	$(c, c^*)(c, c^*)$	self-dual
H_{18}	$C_2 \times C_2$	$(c, c^*)(d, d^*)$	self-dual
H_{19}	$C_2 \times C_2$	$(c, c^*)(c, c^*d^*)$	H_{20}
H_{20}	$C_2 \times C_2$	$(c, c^*)(cd, c^*)$ or $(c, c^*d^*)(d, c^*d^*)$	H_{19}
H_{21}	C_2	$(c, c^*)(c, c^*)(c, c^*)$	self-dual

Note that for some classes, such as H_4 and H_8, we show two choices for g_i and χ_i. To see that the choices for H_8 are isomorphic, use the automorphism ϕ of Γ given by $\phi(c) = c, \phi(d) = c^2d$. The choice $g_8 = d, \chi_8 = c^*d^*$ is natural from Corollary 2.3; the choice $g_8 = c^2d, \chi_8 = c^*$ is mentioned because it corresponds to the listing of the classes in [11]. To confirm the isomorphism for H_4, use the automorphism ϕ of Γ given by $\phi(c) = c^3$. As noted in [11], $H_{10} \cong T \otimes k[C_2 \times C_2]$ where T is the 4-dimensional Sweedler Hopf algebra and so it is obvious that H_{10} is self-dual.

Now we compute some duals before summarizing information on the structure of the duals of the nontrivial liftings. Although Theorem 3.6 does not apply here since r is not an odd integer, the approach is similar.

Example 4.4. Let A be the lifting of H_{14} with matrix $\mathcal{A} = I_2$, i.e. A is the Hopf algebra H_3 of [8, Section 2] or the Hopf algebra $H_1(1)$ from [11, Section 3]. Thus $\Gamma = C_4 =< c >$, $\hat{\Gamma} =< c^* >$ and $V = V_c^{c^{*2}} = kx \oplus ky$ with $xy = -yx$ and $x^2 = y^2 = c^2 - 1$. We describe the dual A^*. In this case, $\Omega = \{1, c^{*2}\}$. Then A^* contains an 8-dimensional Hopf subalgebra $C(\overline{1})$ isomorphic to $\mathcal{B}(W)\#k[\Omega]$ where $W = kw \oplus kz = W^c_{c^{*2}}$. By Lemma 3.1, as an algebra, $A^* \cong \mathcal{B}(W)\#k[\hat{\Gamma}]$. We study the 8-dimensional coalgebra $C(\overline{c^*}) = C(\overline{c^{*3}})$ by finding a matrix subcoalgebra in $C(\overline{c^*})$ explicitly. Note that $c^*(c^2 - 1) = -2$, let i be a primitive 4th root of 1 and let $E = \oplus_{1 \leq i,j \leq 2} ke(i,j)$ with

$$\begin{aligned}
e(1,1) &= c^* - 2iw(c^*, xy);\\
e(1,2) &= w(c^*, x) - iw(c^*, y);\\
e(2,1) &= -2[w(c^{*3}, x) + iw(c^{*3}, y)];\\
e(2,2) &= c^{*3} + 2iw(c^{*3}, xy).
\end{aligned}$$

Then checking that $\{e(1,1), e(1,2), e(2,1), e(2,2)\}$ is a matrix coalgebra basis is a straightforward computation. As noted in Remark 3.7, we have $c^{*2}Ec^{*2} = E$. But $E \neq F = c^{*2}E = Ec^{*2} \cong \mathcal{M}^c(2,k)$. Here F has matrix coalgebra basis $\{f(i,j) | 1 \le i,j \le 2\} = \{e(i,j)c^{*2} | 1 \le i,j \le 2\} = \{c^{*3} - 2iw(c^{*3}, xy), w(c^{*3}, x) - iw(c^{*3}, y), -2[w(c^*, x) + iw(c^*, y)], c^* + 2iw(c^*, xy)\}$. Thus A^* has coradical isomorphic to $k[C_2] \oplus_{i=1}^2 \mathcal{M}^c(2,k)$.

Now let B be the lifting of H_{16} with matrix $\mathcal{A} = I_2$ and note that as algebras $A = B$, i.e. both are generated by C_4, x and y with the same commutation relations. Thus as coalgebras, their duals are isomorphic, i.e., B^* has coradical isomorphic to $k[C_2] \oplus_{i=1}^2 \mathcal{M}^c(2,k)$ and E and F are the same as in A^*. The algebra structure of B^* is determined in Lemma 3.1. □

It remains to describe the dual of the lifting of H_{20} with nondiagonal matrix $\mathcal{A}$. Note that this Hopf algebra A is the Hopf algebra discussed in Example 2.8, i.e. the Drinfel'd double of the Sweedler 4-dimensional Hopf algebra. It is the first lifting we have considered with a nondiagonal matrix $\mathcal{A} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.

Example 4.5. Let A be as above. Then A is generated by $k[\Gamma]$ where $\Gamma =< c > \times < d > \cong C_2 \times C_2$, the $(1,c)$-primitive x and the $(1,cd)$-primitive y. Also $hx = c^*(h)xh$ and $hy = c^*(h)yh$ for all $h \in \Gamma$ where $c^*(c) = d^*(d) = -1$ and $c^*(d) = d^*(c) = +1$. In A, $x^2 = y^2 = 0$, and $xy + yx = d - 1$.

Then $\Omega = \{1, c^*\}$ is the group of grouplikes of A^* and A^* contains a subHopf algebra $C(\overline{1})$ of dimension 8 isomorphic to $\mathcal{B}(W)\#k[\Omega]$ where $W = kw_1 \oplus kw_2$ with $w_1 = w(c^*, x)$ and $w_2 = w(c^*, y)$. Then $w_1 \in W^c_{c^*}$, $w_2 \in W^{cd}_{c^*}$ and $w_1^2 = w_2^2 = 0$.

Note that the standard k-basis we have been using for A is $\{g, gx, gy, gxy | g \in \Gamma\}$ but the set $\{g, g(x+y), g(x-y), g(x+y)(x-y) | g \in \Gamma\}$ is also a basis. Let $x' = x+y, y' = x-y$. Then the set $\{w(\gamma, 1) = \gamma, w(\gamma, x'), w(\gamma, y'), w(\gamma, x'y') | \gamma \in \hat{\Gamma}\}$ is a k-basis for A^* where $w(\gamma, z)(hz) = \gamma(h)$ and $w(\gamma, z)$ is zero on all other basis elements as usual. Note that $2w(\gamma, x') = w(\gamma, x) + w(\gamma, y)$ and also that $w(c^*, x'), w(c^*, y')$ are (ϵ, c^*)-primitive. Thus $C(\overline{d}^*)$ has k-basis $\{w(d^*\gamma, z) | \gamma \in \{1, c^*\}, z \in \{1, x', y', x'y'\}\}$. Note that $x'^2 = -y'^2 = xy + yx = d - 1$. Also $x'y' = yx - xy = -y'x'$. Then $(x'y')^2 = x'(-x'y')y' = (d-1)^2$. The set

$$\begin{aligned}
e(1,1) &= d^* + 2w(d^*, x'y');\\
e(1,2) &= \sqrt{2}w(d^*, x') + \sqrt{2}w(d^*, y')\\
e(2,1) &= -\sqrt{2}w(c^*d^*, x') + \sqrt{2}w(c^*d^*, y');\\
e(2,2) &= c^*d^* - 2w(c^*d^*, x'y')
\end{aligned}$$

is a matrix coalgebra basis for a matrix coalgebra E. A sample computation is:

$$\begin{aligned}
\triangle e(1,1) &= \triangle d^* + 2\triangle w(d^*, x'y') \\
&= [d^* \otimes d^* + (-2)w(d^*, x') \otimes w(c^*d^*, x') + (2)w(d^*, y') \otimes w(c^*d^*, y') \\
&+ \; 4w(d^*, x'y') \otimes w(d^*, x'y')] + 2[d^* \otimes w(d^*, x'y') + w(d^*, x'y') \otimes d^* \\
&+ \; w(d^*, x') \otimes w(c^*d^*, y') - w(d^*, y') \otimes w(c^*d^*.x')] \\
&= d^* \otimes [d^* + 2w(d^*, x'y')] + 2w(d^*, x'y') \otimes [d^* + 2w(d^*, x'y')] \\
&+ \; \sqrt{2}w(d^*, x') \otimes [-\sqrt{2}w(c^*d^*, x') + \sqrt{2}w(c^*d^*, y')] \\
&+ \; \sqrt{2}w(d^*, y') \otimes [\sqrt{2}w(c^*d^*, y') - \sqrt{2}w(c^*d^*, x')] \\
&= e(1,1) \otimes e(1,1) + e(1,2) \otimes e(2,1).
\end{aligned}$$

Note that the first multiplication formula in Lemma 3.1(ii) does not apply to products such as $c^* * w(d^*, x')$ because x' is not the product of skew-primitives. However, the formulas do apply to products such as $c^* * w(d^*, x'y')$ since $\triangle(x'y') = \triangle(yx - xy)$ contains a term $d \otimes x'y'$. Thus we see that $c^*e(1,1)c^* = e(1,1)$ and so $c^*Ec^* = E$. Or we could recall the fact that $c^*Ec^* = E$ from Remark 3.7.

Also $c^*E = Ec^* = F \neq E$ is a matrix coalgebra with matrix coalgebra basis $f(i,j) = e(i,j)c^*$.

Thus we have shown that as a coalgebra $A^* \cong C(\overline{1}) \oplus C(\overline{d^*})$ where $C(\overline{d^*})$ is the direct sum of two 4-dimensional matrix coalgebras, so that corad $(A^*) \cong k[C_2] \oplus C(\overline{d}^*)$. As an algebra, A^* is generated by $\hat{\Gamma}, w_1 = w(c^*, x)$ and $w_2 = w(c^*, y)$ where for $\gamma \in \hat{\Gamma}$, $\gamma w_1 = \gamma(c)w_1\gamma$ and $\gamma w_2 = \gamma(cd)w_2\gamma$; $w_1w_2 = -w_2w_1$; $w_1^2 = w_2^2 = 0$. Alternatively, as an algebra A^* is generated by $\hat{\Gamma}$ and $w_1' = w(c^*, x'), w_2' = w(c^*, y')$ but $\gamma(kw_i')\gamma^{-1} \neq kw_i'$. □

Now we list the duals of the 8 classes of pointed Hopf algebras of dimension 16 which are nontrivial liftings. In the table below, A_i is the nontrivial lifting of the Hopf algebra H_i with matrix $\mathcal{A}_i$, where $D = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $E = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.

A_i	$G(A_i)$	V_i	$\mathcal{A}_i$	A_i^*	$corad(A_i^*)$
A_1	C_8	(c, c^{*4})	I_1	Th. 3.3	$k[C_2] \oplus \oplus_{i=1}^3 \mathcal{M}^c(2,k)$
A_5	$C_4 \times C_2$	(c, c^{*2})	I_1	Th. 3.3	$k[C_2 \times C_2] \oplus \oplus_{i=1}^2 \mathcal{M}^c(2,k)$
A_7	$C_4 \times C_2$	(cd, d^*)	I_1	Th. 3.3	$k[C_2 \times C_2] \oplus \oplus_{i=1}^2 \mathcal{M}^c(2,k)$
$A_{14,1}$	C_4	$(c, c^{*2}), (c, c^{*2})$	D	Cor.3.5	$k[C_2] \oplus \mathcal{M}^c(2,k)$
$A_{14,2}$	C_4	$(c, c^{*2}), (c, c^{*2})$	I_2	Ex. 4.4	$k[C_2] \oplus \oplus_{i=1}^2 \mathcal{M}^c(2,k)$
$A_{16,1}$	C_4	$(c, c^{*2}), (c^3, c^{*2})$	D	Cor.3.5	$k[C_2] \oplus \mathcal{M}^c(2,k)$
$A_{16,2}$	C_4	$(c, c^{*2}), (c^3, c^{*2})$	I_2	Ex. 4.4	$k[C_2] \oplus \oplus_{i=1}^2 \mathcal{M}^c(2,k)$
A_{20}	$C_2 \times C_2$	$(c, c^*), (cd, c^*)$	E	Ex. 4.5	$k[C_2] \oplus \oplus_{i=1}^2 \mathcal{M}^c(2,k)$

4.3. **Duals of an infinite family of dimension 32.** Liftings of quantum linear spaces have provided many counterexamples to Kaplansky's 10th conjecture (see [5], [9], [17]). The examples of smallest dimension are families of Hopf algebras of dimension 32 with coradical $k[\Gamma]$ where Γ is an abelian group of order 8 (see [8], [19]). We compute the duals for the family in [8]. Let $\Gamma =< c >\cong C_8$ and $\hat{\Gamma} =< c^* >$ as usual with $c^*(c) = q$, a primitive 8th root of 1. Let $V = kv_1 \oplus kv_2$ with $v_1 \in V_c^{c^{*4}}, v_2 \in V_{c^5}^{c^{*4}}$ and let $A(\lambda)$ be the lifting of $\mathcal{B}(V)\#k[\Gamma]$ with matrix $\begin{bmatrix} 1 & \lambda \\ \lambda & 1 \end{bmatrix}$. Then $A(\lambda) \ncong A(\lambda')$ unless $\lambda' = \pm\lambda$ by [8], so that the $A(\lambda)$ give an infinite family of nonisomorphic Hopf algebras of dimension 32. Since the original submission of this note, a paper by Etingof and Gelaki has appeared where it is shown that this family contains infinitely many twist equivalence classes [15, Corollary 4.3] , providing a counterexample to a conjecture of Masuoka [25]. Actually Etingof and Gelaki are studying triangular Hopf algebras $A(G, V, u, B)$ in the notation of [15] (see also [2] and [16]) and the $A(\lambda)$ are duals of such Hopf algebras. Here, we compute the coradical of the dual of $A(\lambda)$ using the same type of straightforward computations as in the previous examples.

Example 4.6. For $A = A(\lambda)$ as above, let x, y be the liftings of v_1 and v_2. Then $x^2 = y^2 = c^2 - 1$ and $xy + yx = \lambda(c^6 - 1)$. Then $\Omega = G(A^*) = \{\epsilon, c^{*4}\}$, and A^* contains a subHopf algebra $C(\overline{1})$ of dimension 8 isomorphic to $\mathcal{B}(W)\#k[\Omega]$ where $W = kw_1 \oplus kw_2$, $w_1 = w(c^{*4}, x)$ and $w_2 = w(c^{*4}, y)$. We determine the coalgebra structure of $C(\overline{\gamma}), \gamma \notin \Omega$.

As in Example 4.5, we use a different k-basis for A. Let $x' = x + y, y' = x - y$ and then A has k-basis hz where $h \in \Gamma, z \in \mathcal{Z} = \{1, x', y', x'y'\}$. Then $x'^2 = 2(c^2 - 1) + \lambda(c^6 - 1), y'^2 = 2(c^2 - 1) - \lambda(c^6 - 1)$ and $x'y' = yx - xy = -y'x'$. We use the corresponding basis $\{w(\gamma, z) | \gamma \in \hat{\Gamma}, z \in \mathcal{Z}\}$ for A^*. Let $\gamma \notin \Omega$ and denote $\alpha = \gamma(2(c^2 - 1) + \lambda(c^6 - 1))$, $\beta = \gamma(2(c^2 - 1) - \lambda(c^6 - 1))$.

Case i). Suppose first that $\alpha \neq 0$ and $\beta \neq 0$. Then $E = \oplus_{1 \leq i,j \leq 2} ke(i, j) \cong \mathcal{M}^c(2, k)$ where

$$\begin{aligned} e(1,1) &= \gamma + i\sqrt{\alpha}\sqrt{\beta}w(\gamma, x'y'); \\ e(1,2) &= -\sqrt{\alpha}w(\gamma, x') + i\sqrt{\beta}w(\gamma, y'); \\ e(2,1) &= -\sqrt{\alpha}w(\gamma c^{*4}, x') - i\sqrt{\beta}w(\gamma c^{*4}, y'); \\ e(2,2) &= \gamma c^{*4} - i\sqrt{\alpha}\sqrt{\beta}w(\gamma c^{*4}, x'y'). \end{aligned}$$

(Here $i^2 = -1$.) Note that $E \neq Ec^{*4}$ so $E \neq F = Ec^{*4} \cong \mathcal{M}^c(2, k)$ also and $C(\overline{\gamma}) \cong \mathcal{M}^c(2, k) \oplus \mathcal{M}^c(2, k)$.

Case ii). One of α, β is 0 and the other is nonzero. Suppose that $\alpha \neq 0$ but $\beta = 0$. Then let $E = \oplus_{1 \leq i,j \leq 2} ke(i, j)$ where

$$\begin{aligned} e(1,1) = \gamma \quad &; \quad e(1,2) = -\sqrt{\alpha}w(\gamma, x'); \\ e(2,1) = -\sqrt{\alpha}w(\gamma c^{*4}, x') \quad &; \quad e(2,2) = \gamma c^{*4}; \end{aligned}$$

and it is easy to check that $E \cong \mathcal{M}^c(2, k)$. Here $E = c^{*4}E = Ec^{*4}$. By the same argument as in the proof of Theorem 3.6, any matrix coalgebra basis element $e(i, i)$ must be of the form $\gamma + \zeta w(\gamma, x'y')$. But $\zeta' w(\gamma, x'y') \otimes w(\gamma, x'y')$ does not occur in $\triangle w$ for any basis element w and so $C(\overline{\gamma})$ contains only one matrix coalgebra. It is easy to see that $C(\overline{\gamma})$ has coradical filtration of length 1, i.e. $C(\overline{\gamma}) = C(\overline{\gamma})_1$. The argument for the case $\alpha = 0, \beta \neq 0$ is the same.

Case iii). The case $\alpha = \beta = 0$ cannot occur since then $\gamma(c)^2 = 1$ so that $\gamma \in \{\epsilon, c^{*4}\} = \Omega$.

Now suppose $\gamma = c^*$. Then $\alpha = c^*(c^2 - 1)c^*(2 + \lambda(c^4 + c^2 + 1)) = (q^2 - 1)(2 + q^2\lambda)$ since $q^4 = -1$. So $\alpha = 0$ if and only if $2 + q^2\lambda = 0$. Similarly $\beta = 0$ if and only if $2 - q^2\lambda = 0$. If $\gamma = c^{*3}$ then $\alpha = 0$ if and only if $2 + q^6\lambda = 0$ and $\beta = 0$ if and only if $2 - q^6\lambda = 0$. Thus if $q^2\lambda = \pm 2$, then $C(\overline{c^*})$ and $C(\overline{c^{*3}})$ each have coradical isomorphic to $\mathcal{M}^c(2, k)$. Otherwise both $C(\overline{c^*})$ and $C(\overline{c^{*3}})$ are cosemisimple, isomorphic to $\mathcal{M}^c(2, k) \oplus \mathcal{M}^c(2, k)$.

A similar argument shows that for $\gamma = c^{*2}$, $\alpha = 0$ if and only if $\lambda = -2$ and $\beta = 0$ if and only if $\lambda = 2$. Thus $C(\overline{c^{*2}})$ is cosemisimple unless $\lambda = \pm 2$ and then it has coradical isomorphic to $\mathcal{M}^c(2, k)$.

We summarize in the following table:

λ	$corad(A(\lambda)^*)$
$\pm 2q^2$	$k[C_2] \oplus \oplus_{i=1}^{4} \mathcal{M}^c(2, k)$
± 2	$k[C_2] \oplus \oplus_{i=1}^{5} \mathcal{M}^c(2, k)$
$k - \{\pm 2q^2, \pm 2\}$	$k[C_2] \oplus \oplus_{i=1}^{6} \mathcal{M}^c(2, k)$

Note that in the last row, λ may be 0, and then the result follows from a simpler argument with $\alpha = \beta$. □

Acknowledgement: Many thanks to N. Andruskiewitsch for pointing out more general statements for Theorem 2.2 and Theorem 2.5, and for his many insightful comments, especially on the material in Section 2. Also thanks to S. Dăscălescu and S. Westreich for comments on the original version of Theorem 2.2.

REFERENCES

[1] N. Andruskiewitsch and S. Dăscălescu, On finite quantum groups at -1, Algebr. Represent. Theory 8 (2005), no. 1, 11–34.

[2] N. Andruskiewitsch, P. Etingof, S. Gelaki, Triangular Hopf algebras with the Chevalley property. Michigan Math. J. 49 (2001), no. 2, 277–298.

[3] N. Andruskiewitsch and M. Graña, Braided Hopf algebras over non-abelian finite groups, Boletín de la Acad. Nac. Cs. Córdoba, Argentina 63 (1999), 45-78.

[4] N. Andruskiewitsch and H.-J. Schneider, Pointed Hopf algebras. New directions in Hopf algebras, 1–68, Math. Sci. Res. Inst. Publ., 43, Cambridge Univ. Press, Cambridge, 2002.

[5] N. Andruskiewitsch and H.-J. Schneider, Lifting of quantum linear spaces and pointed Hopf algebras of order p^3, J. Algebra 209 (1998), 659-691.

[6] N. Andruskiewitsch and H.-J. Schneider, Lifting of Nichols algebras of type A_2 and pointed Hopf algebras of order p^4, in “Hopf algebras and quantum groups”, eds. S. Caenepeel and F. van Oystaeyen, M. Dekker, 2000, pp. 1-16.

[7] N. Andruskiewitsch and H.-J. Schneider, Finite quantum groups and Cartan matrices, Adv. Math. 154 (2000), 1–45.

[8] M. Beattie, An isomorphism theorem for Ore extension Hopf algebras, Comm. Algebra 28(2)(2000), 569-584.

[9] M. Beattie, S. Dăscălescu, L. Grünenfelder, Constructing pointed Hopf algebras by Ore extensions, J. Algebra 225, (2000), 743-770.

[10] M. Beattie, S. Dăscălescu and S. Raianu, Lifting of Nichols algebras of type B_2. With an appendix by the authors and Ian Rutherford. Israel J. Math. 132 (2002), 1–28.

[11] S. Caenepeel, S. Dăscălescu, and S. Raianu, Classifying pointed Hopf algebras of dimension 16, Comm. Algebra 28(2)(2000), 541-568.

[12] Călinescu, C.; Dăscălescu, S.; Masuoka, A.; Menini, C. Quantum lines over non-cocommutative cosemisimple Hopf algebras. J. Algebra 273 (2004), no. 2, 753–779.

[13] W. Chin and I. Musson, The coradical filtration for quantized enveloping algebras, J. London Math. Soc. 53 (1996), 50-67.

[14] V. Drinfel'd, Quantum groups, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 798–820, Amer. Math. Soc., Providence, RI, 1987.

[15] P. Etingof and S. Gelaki, On families of triangular Hopf algebras, IMRN **2002:14** (2002), pp. 757-768.

[16] P. Etingof and S. Gelaki, On cotriangular Hopf algebras, Amer. J. Math. 123 (2001) , 699-713.

[17] S. Gelaki, Pointed Hopf algebras and Kaplansky's 10th conjecture, J. Algebra 209 (1998), 635-657.

[18] M. Graña, A freeness theorem for Nichols algebra, J. Algebra, 231 (2000), 235-257.

[19] M. Graña, Pointed Hopf algebras of dimension 32, Comm. Algebra 28 (2000), no. 6, 2935–2976.

[20] Y. Kashina, Classification of semisimple Hopf algebras of dimension 16, J. Algebra 232 (2000), 617-663.

[21] L.H.Kauffman and D.E.Radford, A necessary and sufficient condition for a finite dimensional Drinfel'd double to be a ribbon Hopf algebra, J. Algebra 159 (1993), 98-114.

[22] T.Y. Lam, A First Course in Noncommutative Rings, Springer GTM 131.

[23] G. Lusztig, Introduction to quantum groups, Birkhäuser, 1993.

[24] S. Majid, Foundations of Quantum Group Theory. Cambridge Univ. Press, 1995.

[25] A. Masuoka, Defending the negated Kaplansky conjecture, Proc. Amer. Math. Soc. 129 (2001), 3185-3192.

[26] S. Montgomery, Hopf algebras and their actions on rings, AMS (1993), CBMS **82**.

[27] A. Nenciu, Quasitriangular structures for a class of pointed Hopf algebras constructed by Ore extensions, Comm. Algebra 29 (2001), 1959-1981.

[28] A. Nenciu, Quasitriangular pointed Hopf algebras constructed by Ore extensions, preprint.

[29] W.D. Nichols, Bialgebras of type 1, Comm. Algebra **6** (1978), 1521-1552.

[30] F. Panaite and F. Van Oystaeyen, Quasitriangular structures for some pointed Hopf algebras of dimension 2^n, Comm. Algebra 27(10)(1999), 4929-4942.

[31] D. E. Radford, On the coradical of a finite-dimensional Hopf algebra, Proc. Amer. Math. Soc. 53 (1975), 8-15.

[32] D. E. Radford, Minimal quasitriangular Hopf algebras, J. Algebra 157(1993), 285-315.

[33] D.E. Radford, On Kauffman's knot invariants arising from finite dimensional Hopf algebras, in "Advances in Hopf algebras", pp205-266, J. Bergen and S. Montgomery (eds), Lecture Notes in Pure and Appl. Math 158, Marcel Dekker, New York, 1994.

[34] M.E. Sweedler, Hopf Algebras, Benjamin, New York, 1969.

Department of Mathematics and Computer Science, Mount Allison University, Sackville, N.B., Canada E4L 1E6, email: mbeattie@mta.ca